


5 MONOCLONAL ANTIBODIES TO IFNAR2



FIELD OF THE INVENTION

10 This invention relates to the field of anti-type I interferon receptor antibodies, and more particularly to anti-type I interferon receptor antibodies that block the binding of type I interferons to the second component (IFNAR2) of the type I interferon receptor complex.

BACKGROUND OF THE INVENTION

15 The type I interferons (IFNs) are cytokines which have pleiotropic effects on a wide variety of cell types. IFNs are best known for their anti-viral activity, but they also have anti-bacterial, anti-protozoal, immunomodulatory, and cell-growth regulatory functions. The type 1 interferons include interferon- α (IFN- α) and interferon- β (IFN- β). Human IFN- α (hIFN- α) is a heterogeneous family with at least 23 polypeptides while there is only one IFN- β polypeptide (*J. Interferon Res.*, **13**: 443-444 (1993)). The hIFN- α subtypes show more than 70% amino acid sequence homology, and there is approximately 25% amino acid identity with hIFN- β . The hIFNs- α and hIFN- β share a common receptor.

20 Two components of the hIFN- α receptor complex have recently been cloned. The cDNA for the first hIFN- α receptor (hIFNAR1) encodes a 63 kD receptor protein (reported in Uze *et al.*, *Cell*, **60**: 225-234 (1990)). This receptor undergoes extensive glycosylation which causes it to migrate in gel electrophoresis as a much larger 135 kD protein. The second interferon receptor, hIFNAR2 (hIFN- α β R long), is a 115 kD protein which mediates a functional signaling complex when associated with hIFNAR1 (reported in Domanski *et al.*, *J. Biol. Chem.*, **270**: 21606-21611 (1995)). A variant of IFNAR2, the IFN- α / β receptor (hIFN- α β R short), is a 55 kD protein that can bind to type 1 hIFNs but cannot form a functional complex when associated with hIFNAR1

(reported in Novick *et al.*, *Cell*, **77**: 391-400 (1994)). This IFN- α/β receptor appears to be an alternatively spliced variant of hIFNAR2.

The unprocessed hIFNAR1 expression product is composed of 557 amino acids including an extracellular domain (ECD) of 409 residues, a transmembrane domain of 21 residues, and an intracellular domain of 100 residues as shown in Fig. 5 on page 229 of Uze *et al.*, *supra*. The ECD of IFNAR1 is composed of two domains, domain 1 and domain 2, which are separated by a three-proline motif. There is 19% sequence identity and 50% sequence homology between domains 1 and 2 (Uze *et al.*, *supra*). Each domain (D200) is composed of approximately 200 residues and can be further subdivided into two homologous subdomains (SD100) of approximately 100 amino acids. The unprocessed hIFNAR2 expression product is composed of 515 amino acids, including an extracellular domain (ECD) of 217 residues, a transmembrane domain of 21 residues, and a long cytoplasmic tail of 250 residues as shown in Fig. 1 on page 21608 of Domanski *et al.*, *J. Biol. Chem.*, **37**: 21606-21611 (1995).

Through the use of IFNAR1 gene knockout mice, IFNAR1 has been shown to be essential for the response to all type 1 IFNs (Muller *et al.*, *Science*, **264**: 1918-1921 (1994); Cleary *et al.*, *J. Biol. Chem.*, **269**: 18747-18749 (1994)) and for the mediation of species-specific IFN signal transduction (Constantinescu *et al.*, *Proc. Natl. Acad. Sci. USA*, **91**: 9602-9606 (1994)). However, IFNAR2, not IFNAR1, plays a crucial role in ligand binding (Cohen *et al.*, *Mol. Cell Biol.*, **15**: 4208 (1995)).

Benoit *et al.*, *J. Immunol.*, **150**: 707-716 (1993) reported an anti-IFNAR1 mAb, 64G12, that was found to inhibit the binding of IFN- $\alpha 2$ (IFN- αA) and IFN- αB to Daudi cells and to inhibit the antiviral activity of IFN- $\alpha 2$, IFN- β and IFN- ω (IFN- α_{11}) on Daudi cells. Benoit *et al.* also reported that 64G12 recognizes an epitope present in domain 1 of IFNAR1. Eid and Tovey, *J. Interferon Cytokine Res.*, **15**: 205-211 (1995) reported that 64G12 cannot immunoprecipitate cross-linked IFN- $\alpha 2$ -receptor complexes from Daudi cells.

Colamonici and Domanski, *J. Biol. Chem.*, **268**: 10895-10899 (1993) reported an anti-IFNAR2 mAb (denoted the "IFNAR β 1 mAb") that blocked the binding of IFN- $\alpha 2$ (IFN- αA) to

Daudi cells and U-266 cells and blocked the antiproliferative activity of different type I interferons on Daudi cells using MTT cell proliferation assays.

SUMMARY OF THE INVENTION

In one aspect, the invention provides an anti-IFNAR2 monoclonal antibody which blocks the binding of a first type I interferon to IFNAR2 and does not block the binding of a second type I interferon to IFNAR2.

In another aspect, the invention provides an anti-IFNAR2 monoclonal antibody that competes with an antibody selected from the group consisting of anti-IFNAR2 monoclonal antibodies 1F3, 3B7 and 1D3 for binding to IFNAR2.

In an additional aspect, the invention provides an anti-IFNAR2 monoclonal antibody selected from the group consisting of: (1) an antibody that binds to one or more of amino acid positions 49, 51, 52 and 54 in situ in IFNAR2; (2) an antibody that binds to one or more of amino acid positions 68, 71 and 72 in situ in IFNAR2; (3) an antibody that binds to one or more of amino acid positions 133, 134, 135 and 139 in situ in IFNAR2; (4) an antibody that binds to one or more of amino acid positions 153, 154 and 156 in situ in IFNAR2; (5) an antibody that binds to one or more of amino acid positions 74, 77 and 78 in situ in IFNAR2; and (6) an antibody that binds to one or more of amino acid positions 105 and 109 in situ in IFNAR2.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a graph depicting the ability of anti-IFNAR2 mAbs to inhibit the binding of IFN- α 2/1 to IFNAR2-IgG in an ELISA assay. IFNAR2-IgG was captured onto ELISA wells precoated with goat anti-human IgG Fc. Various concentrations of mAbs and a predetermined suboptimal concentration of biotinylated IFN- α 2/1 (Bio-IFN- α 2/1) were added. The bound Bio-IFN- α 2/1 was detected by the addition of HRP-streptavidin.

Fig. 2 is an autoradiograph depicting the effect of anti-IFNAR2 mAbs on the IGSF3 formation induced by type I IFN. Hela cells were first incubated with mAbs followed by the

addition of IFN- α 8 at a concentration of 20ng/ml. Twenty minutes later, cell lysates were prepared and IGSF complex was detected by electrophoretic mobility shift assay. Anti-IFN- α mAb 9E1 was included as a positive control.

Figs. 3A and 3B are graphs depicting detailed IFNAR2 binding analyses of IFN- α 2/1 and blocking mAb 1F3, respectively. 0.5 nM concentrations of mutant IFNAR2-IgGs were captured onto ELISA wells precoated with goat anti-human IgG Fc. Various concentrations of Bio-IFN- α 2/1 or mAb 1F3 were added. The bound Bio-IFN α 2/1 and mAb 1F3 were detected with HRP-streptavidin and HRP-goat anti-mouse IgG, respectively.

Fig. 4 is a model of hIFNAR2 displaying its protein sequence on the structural backbone of tissue factor. The model shows the location of some residues (in red) important for binding of mAbs to IFNAR2.

Fig. 5 depicts the DNA sequence (SEQ ID NO. 25) and amino acid sequence (SEQ ID NO. 26) of the IFNAR2 ECD-IgG coding insert in pRK5 hIFNAR2-IgG. The DNA sequence encoding the leader peptide amino acid sequence (corresponding to amino acids 1-26 in Fig. 1 on page 21608 of Domanski et al., *J. Biol. Chem.*, **270**: 21606-21611 (1995)) of IFNAR2 is shown as bases 22-99 of SEQ ID NO. 25 in Fig. 5. The leader peptide amino acid sequence is omitted from Fig. 5 in order to present the mature IFNAR2 ECD sequence as amino acids 1-216 of the IFNAR2 ECD-IgG fusion protein sequence (SEQ ID NO. 26). Unless otherwise indicated, the amino acid numbering scheme for IFNAR2 ECD shown in Fig. 5 is used throughout the application.

METHODS OF CARRYING OUT THE INVENTION

A. DEFINITIONS

As used herein, the terms "type I interferon" and "human type I interferon" are defined as all species of native human interferon which fall within the human interferon- α , interferon- ω and interferon- β classes and which bind to a common cellular receptor. Natural human interferon- α comprises 23 or more closely related proteins encoded by distinct genes with a high

degree of structural homology (Weissmann and Weber, *Prog. Nucl. Acid. Res. Mol. Biol.*, **33**: 251 (1986); *J. Interferon Res.*, **13**: 443-444 (1993)). The human IFN- α locus comprises two subfamilies. The first subfamily consists of at least 14 functional, non-allelic genes, including genes encoding IFN- α A (IFN- α 2), IFN- α B (IFN- α 8), IFN- α C (IFN- α 10), IFN- α D (IFN- α 1), IFN- α E (IFN- α 22), IFN- α F (IFN- α 21), IFN- α G (IFN- α 5), and IFN- α H (IFN- α 14), and pseudogenes having at least 80% homology. The second subfamily, α_{II} or ω , contains at least 5 pseudogenes and 1 functional gene (denoted herein as "IFN- α_{II} 1" or "IFN- ω ") which exhibits 70% homology with the IFN- α genes (Weissmann and Weber (1986)). The human IFN- β is encoded by a single copy gene.

As used herein, the terms "first human interferon- α (hIFN- α) receptor", "IFN- α R", "hIFNAR1", "IFNAR1", and "Uze chain" are defined as the 557 amino acid receptor protein cloned by Uze *et al.*, Cell, **60**: 225-234 (1990), including an extracellular domain of 409 residues, a transmembrane domain of 21 residues, and an intracellular domain of 100 residues, as shown in Fig. 5 on page 229 of Uze *et al.* Also encompassed by the foregoing terms are fragments of IFNAR1 that contain the extracellular domain (ECD) (or fragments of the ECD) of IFNAR1.

As used herein, the terms "second human interferon- α (hIFN- α) receptor", "IFN- α β R", "hIFNAR2", "IFNAR2", and "Novick chain" are defined as the 515 amino acid receptor protein cloned by Domanski *et al.*, J. Biol. Chem., **37**: 21606-21611 (1995), including an extracellular domain of 217 residues, a transmembrane domain of 21 residues, and an intracellular domain of 250 residues, as shown in Fig. 1 on page 21608 of Domanski *et al.* Also encompassed by the foregoing terms are fragments of IFNAR2 that contain the extracellular domain (ECD) (or fragments of the ECD) of IFNAR2, and soluble forms of IFNAR2, such as IFNAR2 ECD fused to an immunoglobulin sequence, e.g. IFNAR2 ECD-IgG as described in the Example below.

As used herein, the term "anti-IFNAR2 antibody" is defined as an antibody that is capable of binding to IFNAR2.

As used herein, an anti-IFNAR2 antibody with the property or capability of "blocking the binding of a type I interferon to IFNAR2" is defined as an anti-IFNAR2 antibody capable of binding to IFNAR2 such that the ability of IFNAR2 to bind to one or more type I interferons is impaired or eliminated. An anti-IFNAR2 antibody candidate can be tested for such activity, for example, by adsorbing anti-IFNAR2 antibody to immobilized IFNAR2 followed by subjecting the adsorbed antibody to elution with an excess of a selected type I interferon. If an eluent comprising an excess of the selected type I interferon produces an eluate containing a greater concentration of the candidate antibody than the concentration of candidate antibody present in an eluate produced by a "blank" eluent (the same eluent containing no type I interferon) in a control elution, as determined by, e.g., radioimmunoassays performed on the respective eluates with radiolabelled, soluble IFNAR2, then the candidate antibody competes with the selected type I interferon for binding to IFNAR2. In some embodiments, the anti-IFNAR2 antibody of the invention competes with a selected type I interferon for binding to IFNAR2 and accordingly impairs or eliminates the binding of the selected type I interferon to IFNAR2.

"Polymerase chain reaction" or "PCR" refers to a procedure or technique in which minute amounts of a specific piece of nucleic acid, RNA and/or DNA, are amplified as described in U.S. Patent No. 4,683,195 issued 28 July 1987. Generally, sequence information from the ends of the region of interest or beyond needs to be available, such that oligonucleotide primers can be designed; these primers will be identical or similar in sequence to opposite strands of the template to be amplified. The 5' terminal nucleotides of the two primers can coincide with the ends of the amplified material. PCR can be used to amplify specific RNA sequences, specific DNA sequences from total genomic DNA, and cDNA transcribed from total cellular RNA, bacteriophage or plasmid sequences, etc. See generally Mullis *et al.*, *Cold Spring Harbor Symp. Quant. Biol.* **51**:263 (1987); Erlich, ed., *PCR Technology* (Stockton Press, NY, 1989). As used herein, PCR is considered to be one, but not the only, example of a nucleic acid polymerase reaction method for amplifying a nucleic acid test sample comprising the use of a known nucleic

acid as a primer and a nucleic acid polymerase to amplify or generate a specific piece of nucleic acid.

"Antibodies" (Abs) and "immunoglobulins" (Igs) are glycoproteins having the same structural characteristics. While antibodies exhibit binding specificity to a specific antigen, immunoglobulins include both antibodies and other antibody-like molecules which lack antigen specificity. Polypeptides of the latter kind are, for example, produced at low levels by the lymph system and at increased levels by myelomas.

"Native antibodies and immunoglobulins" are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies between the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (VH) followed by a number of constant domains. Each light chain has a variable domain at one end (VL) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light- and heavy-chain variable domains (Clothia *et al.*, *J. Mol. Biol.* **186**:651 (1985); Novotny and Haber, *Proc. Natl. Acad. Sci. U.S.A.* **82**:4592 (1985)).

The term "variable" refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called complementarity-determining regions (CDRs) or hypervariable regions both in the light-chain and the heavy-chain variable domains. The more highly conserved portions of variable domains are called the framework (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a β -sheet configuration, connected by three CDRs,

which form loops connecting, and in some cases forming part of, the β -sheet structure. The CDRs in each chain are held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, Fifth Edition, National Institute of Health, Bethesda, MD (1991)). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an $F(ab')_2$ fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. In a two-chain Fv species, this region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. In a single-chain Fv species, one heavy- and one light-chain variable domain can be covalently linked by a flexible peptide linker such that the light and heavy chains can associate in a "dimeric" structure analogous to that in a two-chain Fv species. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the VH-VL dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. $F(ab')_2$ antibody

fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

5 Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these can be further divided into subclasses (isotypes), e.g., IgG₁, IgG₂, IgG₃, IgG₄, IgA₁, and IgA₂. The heavy-chain constant domains that correspond to the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

The term "antibody" specifically covers monoclonal antibodies, including antibody fragment clones.

"Antibody fragments" comprise a portion of an intact antibody, generally the antigen binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; single-chain antibody molecules, including single-chain Fv (scFv) molecules; and multispecific antibodies formed from antibody fragments.

The term "monoclonal antibody" as used herein refers to an antibody (or antibody fragment) obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single
20
25 determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma culture, uncontaminated by other

immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (see, *e.g.*, U.S. Patent No. 4,816,567). The "monoclonal antibodies" also include clones of antigen-recognition and binding-site containing antibody fragments (Fv clones) isolated from phage antibody libraries using the techniques described in Clackson *et al.*, *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991), for example.

The monoclonal antibodies herein specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567 to Cabilly *et al.*; Morrison *et al.*, *Proc. Natl. Acad. Sci. USA*, 81:6851-6855 (1984)).

"Humanized" forms of non-human (*e.g.*, murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementarity-determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues which are found neither in the recipient antibody nor in the imported CDR or

framework sequences. These modifications are made to further refine and optimize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, *Nature*, **321**:522-525 (1986); Reichmann *et al.*, *Nature*, **332**:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.*, **2**:593-596 (1992). The humanized antibody includes a Primatized™ antibody wherein the antigen-binding region of the antibody is derived from an antibody produced by immunizing macaque monkeys with the antigen of interest.

"Single-chain Fv" or "scFv" antibody fragments comprise the VH and VL domains of antibody, wherein these domains are present in a single polypeptide chain. Generally, the scFv polypeptide further comprises a polypeptide linker between the VH and VL domains which enables the scFv to form the desired structure for antigen binding. For a review of scFv see Pluckthun, in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (VH) connected to a light-chain variable domain (VL) in the same polypeptide chain (VH - VL). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, **90**:6444-6448 (1993).

An "isolated" antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the

antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody *in situ* within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

"Treatment" refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment include those already with the disorder as well as those in which the disorder is to be prevented.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, *etc.* Preferably, the mammal is human.

As used herein, the terms "each member of the group consisting of" and "each of" are synonymous.

As used herein, the terms "any member of the group consisting of" and "any of" are synonymous.

B. GENERAL METHODS

In general, the invention provides anti-IFNAR2 antibodies that are useful for treatment of immune-mediated disorders in which a partial or total blockade of type I interferon activity is desired. In one embodiment, the anti-IFNAR2 antibodies of the invention are used to treat autoimmune disorders, such as type I and type II diabetes, systemic lupus erythematosus (SLE), and rheumatoid arthritis. In another embodiment, the anti-IFNAR2 antibodies provided herein are used to treat graft rejection or graft versus host disease. The unique properties of the anti-

IFNAR2 antibodies of the invention make them particularly useful for effecting target levels of immunosuppression in a patient. For patients requiring acute intervention, the anti-IFNAR2 antibodies provided herein which cause broad spectrum ablation of type I interferon activity can be used to effect the largest possible compromise of an undesired immune response. For patients requiring maintenance immunosuppression, the anti-IFNAR2 antibodies provided herein which
5 block one or more (but not all) species of type I interferon can be used to effect partial compromise of the patient's immune system in order to reduce the risk of undesirable immune responses while leaving some components of the patient's type I interferon-mediated immunity intact in order to avoid infection.

In another aspect, the anti-IFNAR2 antibodies of the invention find utility as reagents for detection and isolation of IFNAR2, such as detection of IFNAR2 expression in various cell types and tissues, including the determination of IFNAR2 receptor density and distribution in cell populations, and cell sorting based on IFNAR2 expression. In yet another aspect, the present anti-IFNAR2 antibodies are useful for the development of IFNAR2 antagonists with type I interferon blocking activity patterns similar to those of the subject antibodies. The anti-IFNAR2 antibodies of the invention can be used in IFNAR2 signal transduction assays to screen for small molecule antagonists of IFNAR2 which will exhibit similar pharmacological effects in blocking the binding of type I interferons to IFNAR2.

I. Methods of Making Synthetic Anti-IFNAR2 Fv Clones

20 The anti-IFNAR2 antibodies of the invention can be made by using combinatorial libraries to screen for synthetic antibody clones with the desired activity or activities. In principle, synthetic antibody clones are selected by screening phage libraries containing phage that display various fragments of antibody variable region (Fv) fused to phage coat protein. Such phage libraries are panned by affinity chromatography against the desired antigen. Clones
25 expressing Fv fragments capable of binding to the desired antigen are adsorbed to the antigen and thus separated from the non-binding clones in the library. The binding clones are then eluted

from the antigen, and can be further enriched by additional cycles of antigen adsorption/elution. Any of the anti-IFNAR2 antibodies of the invention can be obtained by designing a suitable antigen screening procedure to select for the phage clone of interest followed by construction of a full length anti-IFNAR2 antibody clone using the Fv sequences from the phage clone of interest and suitable constant region (Fc) sequences described in Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, Fifth Edition, NIH Publication 91-3242, Bethesda MD (1991), vols. 1-3.

1. Construction of Phage Libraries

The antigen-binding domain of an antibody is formed from two variable (V) regions of about 110 amino acids, one each from the light (VL) and heavy (VH) chains, that both present three hypervariable loops or complementarity-determining regions (CDRs). Variable domains can be displayed functionally on phage, either as single-chain Fv (scFv) fragments, in which VH and VL are covalently linked through a short, flexible peptide, or as Fab fragments, in which they are each fused to a constant domain and interact non-covalently, as described in Winter *et al.*, *Ann. Rev. Immunol.*, **12**: 433-455 (1994). As used herein, scFv encoding phage clones and Fab encoding phage clones are collectively referred to as "Fv phage clones" or "Fv clones".

The naive repertoire of an animal (the repertoire before antigen challenge) provides it with antibodies that can bind with moderate affinity (K_d^{-1} of about 10^6 to 10^7 M⁻¹) to essentially any non-self molecule. The sequence diversity of antibody binding sites is not encoded directly in the germline but is assembled in a combinatorial manner from V gene segments. In human heavy chains, the first two hypervariable loops (H1 and H2) are drawn from less than 50 VH gene segments, which are combined with D segments and JH segments to create the third hypervariable loop (H3). In human light chains, the first two hypervariable loops (L1 and L2) and much of the third (L3) are drawn from less than approximately 30 V λ and less than approximately 30 V κ segments to complete the third hypervariable loop (L3).

Each combinatorial rearrangement of V-gene segments in stem cells gives rise to a B cell that expresses a single VH-VL combination. Immunizations triggers any B cells making a

combination that binds the immunogen to proliferate (clonal expansion) and to secrete the corresponding antibody. These naive antibodies are then matured to high affinity ($K_d^{-1} \geq 10^9 \text{ M}^{-1}$) by a process of mutagenesis and selection known as affinity maturation. It is after this point that cells are normally removed to prepare hybridomas and generate high-affinity monoclonal antibodies.

At three stages of this process, repertoires of VH and VL genes can be separately cloned by polymerase chain reaction (PCR) and recombined randomly in phage libraries, which can then be searched for antigen-binding clones as described in Winter *et al.*, *Ann. Rev. Immunol.*, **12**: 433-455 (1994). Libraries from immunized sources provide high-affinity antibodies to the immunogen without the requirement of constructing hybridomas. Alternatively, the naive repertoire can be cloned to provide a single source of human antibodies to a wide range of non-self and also self antigens without any immunization as described by Griffiths *et al.*, *EMBO J.*, **12**: 725-734 (1993). Finally, naive libraries can also be made synthetically by cloning the unrearranged V-gene segments from stem cells, and using PCR primers containing random sequence to encode the highly variable CDR3 regions and to accomplish rearrangement in vitro as described by Hoogenboom and Winter, *J. Mol. Biol.*, **227**: 381-388 (1992).

Phage display mimics the B cell. Filamentous phage is used to display antibody fragments by fusion to the minor coat protein pIII. The antibody fragments can be displayed as single chain Fv fragments, in which VH and VL domains are connected on the same polypeptide chain by a flexible polypeptide spacer, e.g. as described by Marks *et al.*, *J. Mol. Biol.*, **222**: 581-597 (1991), or as Fab fragments, in which one chain is fused to pIII and the other is secreted into the bacterial host cell periplasm where assembly of a Fab-coat protein structure which becomes displayed on the phage surface by displacing some of the wild type coat proteins, e.g. as described in Hoogenboom *et al.*, *Nucl. Acids Res.*, **19**: 4133-4137 (1991). When antibody fragments are fused to the N-terminus of pIII, the phage is infective. However, if the N-terminal domain of pIII is excised and fusions made to the second domain, the phage is not infective, and wild type pIII must be provided by helper phage.

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

The pIII fusion and other proteins of the phage can be encoded entirely within the same phage replicon, or on different replicons. When two replicons are used, the pIII fusion is encoded on a phagemid, a plasmid containing a phage origin of replication. Phagemids can be packaged into phage particles by "rescue" with a helper phage such as M13K07 that provides all the phage proteins, including pIII, but due to a defective origin is itself poorly packaged in competitions with the phagemids as described in Vieira and Messing, *Meth. Enzymol.*, **153**: 3-11 (1987). In a preferred method, the phage display system is designed such that the recombinant phage can be grown in host cells under conditions permitting no more than a minor amount of phage particles to display more than one copy of the Fv-coat protein fusion on the surface of the particle as described in Bass *et al.*, *Proteins*, **8**: 309-314 (1990) and in WO 92/09690 (PCT/US91/09133 published June 11, 1992).

In general, nucleic acids encoding antibody gene fragments are obtained from immune cells harvested from humans or animals. If a library biased in favor of anti-IFNAR2 clones is desired, the subject is immunized with IFNAR2 to generate an antibody response, and spleen cells and/or circulating B cells other peripheral blood lymphocytes (PBLs) are recovered for library construction. In a preferred embodiment, a human antibody gene fragment library biased in favor of anti-human IFNAR2 clones is obtained by generating an anti-human IFNAR2 antibody response in transgenic mice carrying a functional human immunoglobulin gene array (and lacking a functional endogenous antibody production system) such that IFNAR2 immunization gives rise to B cells producing human antibodies against IFNAR2. The generation of human antibody-producing transgenic mice is described in Section B(III)(b) below.

Additional enrichment for anti-IFNAR2 reactive cell populations can be obtained by using a suitable screening procedure to isolate B cells expressing IFNAR2-specific membrane bound antibody, e.g., by cell separation with IFNAR2 affinity chromatography or adsorption of cells to fluorochrome-labelled IFNAR2 followed by flow-activated cell sorting (FACS).

Alternatively, the use of spleen cells and/or B cells or other PBLs from an unimmunized donor provides a better representation of the possible antibody repertoire, and also permits the

construction of an antibody library using any animal (human or non-human) species in which IFNAR2 is not antigenic. For libraries incorporating in vitro antibody gene construction, stem cells are harvested from the subject to provide nucleic acids encoding unrearranged antibody gene segments. The immune cells of interest can be obtained from a variety of animal species, such as human, mouse, rat, lagomorpha, luprine, canine, feline, porcine, bovine, equine, and avian species, etc.

Nucleic acid encoding antibody variable gene segments (including VH and VL segments) are recovered from the cells of interest and amplified. In the case of rearranged VH and VL gene libraries, the desired DNA can be obtained by isolating genomic DNA or mRNA from lymphocytes followed by polymerase chain reaction (PCR) with primers matching the 5' and 3' ends of rearranged VH and VL genes as described in Orlandi *et al.*, *Proc. Natl. Acad. Sci. (USA)*, **86**: 3833-3837 (1989), thereby making diverse V gene repertoires for expression. The V genes can be amplified from cDNA and genomic DNA, with back primers at the 5' end of the exon encoding the mature V-domain and forward primers based within the J-segment as described in Orlandi *et al.* (1989) and in Ward *et al.*, *Nature*, **341**: 544-546 (1989). However, for amplifying from cDNA, back primers can also be based in the leader exon as described in Jones *et al.*, *Biotechnol.*, **9**: 88-89 (1991), and forward primers within the constant region as described in Sastry *et al.*, *Proc. Natl. Acad. Sci. (USA)*, **86**: 5728-5732 (1989). To maximize complementarity, degeneracy can be incorporated in the primers as described in Orlandi *et al.* (1989) or Sastry *et al.* (1989). Preferably, the library diversity is maximized by using PCR primers targeted to each V-gene family in order to amplify all available VH and VL arrangements present in the immune cell nucleic acid sample, e.g. as described in the method of Marks *et al.*, *J. Mol. Biol.*, **222**: 581-597 (1991) or as described in the method of Orum *et al.*, *Nucleic Acids Res.*, **21**: 4491-4498 (1993). For cloning of the amplified DNA into expression vectors, rare restriction sites can be introduced within the PCR primer as a tag at one end as described in Orlandi *et al.* (1989), or by further PCR amplification with a tagged primer as described in Clackson *et al.*, *Nature*, **352**: 624-628 (1991).

Repertoires of synthetically rearranged V genes can be derived in vitro from V gene segments. Most of the human VH-gene segments have been cloned and sequenced (reported in Tomlinson *et al.*, *J. Mol. Biol.*, **227**: 776-798 (1992)), and mapped (reported in Matsuda *et al.*, *Nature Genet.*, **3**: 88-94 (1993)); these cloned segments (including all the major conformations of the H1 and H2 loop) can be used to generate diverse VH gene repertoires with PCR primers encoding H3 loops of diverse sequence and length as described in Hoogenboom and Winter, *J. Mol. Biol.*, **227**: 381-388 (1992). VH repertoires can also be made with all the sequence diversity focussed in a long H3 loop of a single length as described in Barbas *et al.*, *Proc. Natl. Acad. Sci. USA*, **89**: 4457-4461 (1992). Human V κ and V λ segments have been cloned and sequenced (reported in Williams and Winter, *Eur. J. Immunol.*, **23**: 1456-1461 (1993)) and can be used to make synthetic light chain repertoires. Synthetic V gene repertoires, based on a range of VH and VL folds, and L3 and H3 lengths, will encode antibodies of considerable structural diversity. Following amplification of V-gene encoding DNAs, germline V-gene segments can be rearranged in vitro according to the methods of Hoogenboom and Winter, *J. Mol. Biol.*, **227**: 381-388 (1992).

Repertoires of antibody fragments can be constructed by combining VH and VL gene repertoires together in several ways. Each repertoire can be created in different vectors, and the vectors recombined in vitro, e.g., as described in Hogrefe *et al.*, *Gene*, **128**: 119-126 (1993), or in vivo by combinatorial infection, e.g., the loxP system described in Waterhouse *et al.*, *Nucl. Acids Res.*, **21**: 2265-2266 (1993). The *in vivo* recombination approach exploits the two-chain nature of Fab fragments to overcome the limit on library size imposed by *E. coli* transformation efficiency. Naive VH and VL repertoires are cloned separately, one into a phagemid and the other into a phage vector. The two libraries are then combined by phage infection of phagemid-containing bacteria so that each cell contains a different combination and the library size is limited only by the number of cells present (about 10^{12} clones). Both vectors contain in vivo recombination signals so that the VH and VL genes are recombined onto a single replicon and

are co-packaged into phage virions. These huge libraries provide large numbers of diverse antibodies of good affinity (K_d^{-1} of about 10^{-8} M).

Alternatively, the repertoires may be cloned sequentially into the same vector, e.g. as described in Barbas *et al.*, *Proc. Natl. Acad. Sci. USA*, **88**: 7978-7982 (1991), or assembled together by PCR and then cloned, e.g. as described in Clackson *et al.*, *Nature*, **352**: 624-628 (1991). PCR assembly can also be used to join VH and VL DNAs with DNA encoding a flexible peptide spacer to form single chain Fv (scFv) repertoires. In yet another technique, "in cell PCR assembly" is used to combine VH and VL genes within lymphocytes by PCR and then clone repertoires of linked genes as described in Embleton *et al.*, *Nucl. Acids Res.*, **20**: 3831-3837 (1992).

The antibodies produced by naive libraries (either natural or synthetic) can be of moderate affinity (K_d^{-1} of about 10^6 to 10^7 M⁻¹), but affinity maturation can also be mimicked *in vitro* by constructing and reselecting from secondary libraries as described in Winter *et al.* (1994), *supra*. For example, mutation can be introduced at random in vitro by using error-prone polymerase (reported in Leung *et al.*, *Technique*, **1**: 11-15 (1989)) in the method of Hawkins *et al.*, *J. Mol. Biol.*, **226**: 889-896 (1992) or in the method of Gram *et al.*, *Proc. Natl. Acad. Sci. USA*, **89**: 3576-3580 (1992). Additionally, affinity maturation can be performed by randomly mutating one or more CDRs, e.g. using PCR with primers carrying random sequence spanning the CDR of interest, in selected individual Fv clones and screening for higher affinity clones. WO 9607754 (published 14 March 1996) described a method for inducing mutagenesis in a complementarity determining region of an immunoglobulin light chain to create a library of light chain genes. Another effective approach is to recombine the VH or VL domains selected by phage display with repertoires of naturally occurring V domain variants obtained from unimmunized donors and screen for higher affinity in several rounds of chain reshuffling as described in Marks *et al.*, *Biotechnol.*, **10**: 779-783 (1992). This technique allows the production of antibodies and antibody fragments with affinities in the 10^{-9} M range.

2. Panning Phage Display Libraries for Anti-IFNAR2 Clones

a. Synthesis of IFNAR2 and IFNAR2 Ligands

Nucleic acid sequence encoding the IFNAR2s used herein can be designed using the amino acid sequence of the desired region of IFNAR2, e.g. the extracellular domain spanning amino acids 27 to 243 of Fig. 7 on page 395 of Novick et al., Cell, **77**: 391-400 (1994). Alternatively, the cDNA sequence of Fig. 7 on page 395 of Novick et al., supra, can be used. In addition, nucleic acid encoding an immunoglobulin G (IgG)-IFNAR2 extracellular domain fusion protein can be obtained from the amino acid or cDNA sequence shown in Fig. 5 below. Likewise, nucleic acid sequence encoding the human type I interferons used herein can be designed using published amino acid and nucleic acid sequences, e.g. see the *J. Interferon Res.*, **13**: 443-444 (1993) compilation of references containing genomic and cDNA sequences for various type I interferons, and the references cited therein. For the IFN- α A, IFN- α B, IFN- α C, IFN- α D, IFN- α E, IFN- α F, IFN- α G, and IFN- α H amino acid sequences or cDNA sequences, see Figs. 3 and 4 on pages 23-24 of Goeddel et al., *Nature*, **290**: 20-26 (1981). For cDNA encoding the amino acid sequence of IFN- α_{11} (IFN- ω), see Capon et al., *Mol. Cell. Biol.*, **5**: 768-779 (1985) and Hauptmann and Swetly, *Nucleic Acids Res.*, **13**: 4739-4749 (1985). For cDNA encoding the amino acid sequence of IFN- β , see Taniguchi et al., *Proc. Jpn. Acad. Ser. B*, **55**: 464-469 (1979); Taniguchi et al., *Gene*, **10**: 11-15 (1980); and U.S. Pat. No. 5,460,811 to Goeddel and Crea. For cDNA encoding the amino acid sequence of IFN- α 7, see Cohen et al., *Dev. Biol. Standard*, **60**: 111-122 (1985). DNAs encoding the IFNAR2s or type I interferons of interest can be prepared by a variety of methods known in the art. These methods include, but are not limited to, chemical synthesis by any of the methods described in Engels et al., *Angew. Chem. Int. Ed. Engl.*, **28**: 716-734 (1989), such as the triester, phosphite, phosphoramidite and H-phosphonate methods. In one embodiment, codons preferred by the expression host cell are used in the design of the IFNAR2 or type I interferon-encoding DNA. Alternatively, DNA encoding the IFNAR2 or type I interferon can be isolated from a genomic or cDNA library.

For production of the mutant IFNAR2s used herein, DNA sequence encoding wild type IFNAR2 can be altered to encode the desired IFNAR2 mutant by using recombinant DNA techniques, such as site specific mutagenesis (Kunkel *et al.*, *Methods Enzymol.* **204**:125-139 (1991); Carter, P., *et al.*, *Nucl. Acids. Res.* **13**:4331 (1986); Zoller, M. J. *et al.*, *Nucl. Acids Res.* **10**:6487 (1982)), cassette mutagenesis (Wells, J. A., *et al.*, *Gene* **34**:315 (1985)), restriction selection mutagenesis (Wells, J. A., *et al.*, *Philos. Trans, R. Soc. London SerA* **317**: 415 (1986)), and the like.

Following construction of the DNA molecule encoding the IFNAR2 or type I interferon of interest, the DNA molecule is operably linked to an expression control sequence in an expression vector, such as a plasmid, wherein the control sequence is recognized by a host cell transformed with the vector. In general, plasmid vectors contain replication and control sequences which are derived from species compatible with the host cell. The vector ordinarily carries a replication site, as well as sequences which encode proteins that are capable of providing phenotypic selection in transformed cells.

For expression in prokaryotic hosts, suitable vectors include pBR322 (ATCC No. 37,017), pHGH107 (ATCC No. 40,011), pBO475, pS0132, pRIT5, any vector in the pRIT20 or pRIT30 series (Nilsson and Abrahmsen, *Meth. Enzymol.*, **185**: 144-161 (1990)), pRIT2T, pKK233-2, pDR540 and pPL-lambda. Prokaryotic host cells containing the expression vectors suitable for use herein include *E. coli* K12 strain 294 (ATCC NO. 31446), *E. coli* strain JM101 (Messing *et al.*, *Nucl. Acid Res.*, **9**: 309 (1981)), *E. coli* strain B, *E. coli* strain χ 1776 (ATCC No. 31537), *E. coli* c600 (Appleyard, *Genetics*, **39**: 440 (1954)), *E. coli* W3110 (F-, gamma-, prototrophic, ATCC No. 27325), *E. coli* strain 27C7 (W3110, *tonA*, *phoA* *E15*, (*argF-lac*)169, *ptr3*, *degP41*, *ompT*, *kan^r*) (U.S. Patent No. 5,288,931, ATCC No. 55,244), *Bacillus subtilis*, *Salmonella typhimurium*, *Serratia marcesans*, and *Pseudomonas* species.

In addition to prokaryotes, eukaryotic organisms, such as yeasts, or cells derived from multicellular organisms can be used as host cells. For expression in yeast host cells, such as common baker's yeast or *Saccharomyces cerevisiae*, suitable vectors include episomally

replicating vectors based on the 2-micron plasmid, integration vectors, and yeast artificial chromosome (YAC) vectors. For expression in insect host cells, such as Sf9 cells, suitable vectors include baculoviral vectors. For expression in plant host cells, particularly dicotyledonous plant hosts, such as tobacco, suitable expression vectors include vectors derived from the Ti plasmid of *Agrobacterium tumefaciens*.

5 However, interest has been greatest in vertebrate host cells. Examples of useful mammalian host cells include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, *J. Gen Virol.*, **36**: 59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA*, **77**: 4216 (1980)); mouse sertoli cells (TM4, Mather, *Biol. Reprod.*, **23**: 243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather *et al.*, *Annals N.Y. Acad. Sci.*, **383**: 44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma cell line (Hep G2). For expression in mammalian host cells, useful vectors include vectors derived from SV40, vectors derived from cytomegalovirus such as the pRK vectors, including pRK5 and pRK7 (Suva *et al.*, *Science*, **237**: 893-896 (1987), EP 307,247 (3/15/89), EP 278,776 (8/17/88)) vectors derived from vaccinia viruses or other pox viruses, and retroviral vectors such as vectors derived from Moloney's murine leukemia virus (MoMLV).

Optionally, the DNA encoding the IFNAR2 or type I interferon of interest is operably linked to a secretory leader sequence resulting in secretion of the expression product by the host cell into the culture medium. Examples of secretory leader sequences include stII, ecotin, lamB, herpes GD, lpp, alkaline phosphatase, invertase, and alpha factor. Also suitable for use herein is the 36 amino acid leader sequence of protein A (Abrahmsen *et al.*, *EMBO J.*, **4**: 3901 (1985)).

Host cells are transfected and preferably transformed with the above-described expression or cloning vectors of this invention and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

Transfection refers to the taking up of an expression vector by a host cell whether or not any coding sequences are in fact expressed. Numerous methods of transfection are known to the ordinarily skilled artisan, for example, CaPO₄ precipitation and electroporation. Successful transfection is generally recognized when any indication of the operation of this vector occurs within the host cell.

Transformation means introducing DNA into an organism so that the DNA is replicable, either as an extrachromosomal element or by chromosomal integrant. Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in section 1.82 of Sambrook *et al.*, *Molecular Cloning* (2nd ed.), Cold Spring Harbor Laboratory, NY (1989), is generally used for prokaryotes or other cells that contain substantial cell-wall barriers. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw *et al.*, *Gene*, **23**: 315 (1983) and WO 89/05859 published 29 June 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method described in sections 16.30-16.37 of Sambrook *et al.*, *supra*, is preferred. General aspects of mammalian cell host system transformations have been described by Axel in U.S. 4,399,216 issued 16 August 1983. Transformations into yeast are typically carried out according to the method of Van Solingen *et al.*, *J. Bact.*, **130**: 946 (1977) and Hsiao *et al.*, *Proc. Natl. Acad. Sci. (USA)*, **76**: 3829 (1979). However, other methods for introducing DNA into cells such as by nuclear injection, electroporation, or by protoplast fusion may also be used.

Prokaryotic host cells used to produce the IFNAR2 or type I interferon of interest can be cultured as described generally in Sambrook *et al.*, *supra*.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
2233
2234
2235
2236
2237

membrane bound peptide can be recovered from the membrane fraction by solubilization with detergents. The crude peptide extract can then be further purified by suitable procedures such as fractionation on immunoaffinity or ion-exchange columns; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; hydrophobic affinity resins and ligand affinity using IFNAR2 (for type I interferon purification) or type I interferons or anti-IFNAR2 antibodies (for IFNAR2 purification) immobilized on a matrix.

Many of the human type I interferons used herein can be obtained from commercial sources, e.g. human IFN- β is available from Sigma (St. Louis, MO).

b. Immobilization of IFNAR2

The purified IFNAR2 can be attached to a suitable matrix such as agarose beads, acrylamide beads, glass beads, cellulose, various acrylic copolymers, hydroxyl methacrylate gels, polyacrylic and polymethacrylic copolymers, nylon, neutral and ionic carriers, and the like, for use in the affinity chromatographic separation of phage display clones.

Attachment of the IFNAR2 protein to the matrix can be accomplished by the methods described in *Methods in Enzymology*, vol. 44 (1976). A commonly employed technique for attaching protein ligands to polysaccharide matrices, e.g. agarose, dextran or cellulose, involves activation of the carrier with cyanogen halides and subsequent coupling of the peptide ligand's primary aliphatic or aromatic amines to the activated matrix.

Alternatively, IFNAR2 can be used to coat the wells of adsorption plates, expressed on host cells affixed to adsorption plates or used in cell sorting, or conjugated to biotin for capture with streptavidin-coated beads, or used in any other art-known method for panning phage display libraries.

c. Panning Procedures

The phage library samples are contacted with immobilized IFNAR2 under conditions suitable for binding of at least a portion of the phage particles with the adsorbent. Normally, the conditions, including pH, ionic strength, temperature and the like are selected to mimic physiological conditions. The phages bound to the solid phase are washed and then eluted by acid, e.g. as described in Barbas *et al.*, *Proc. Natl. Acad. Sci USA*, **88**: 7978-7982 (1991), or by alkali, e.g. as described in Marks *et al.*, *J. Mol. Biol.*, **222**: 581-597 (1991), or by IFNAR2 antigen or type I interferon ligand competition, e.g. in a procedure similar to the antigen competition method of Clackson *et al.*, *Nature*, **352**: 624-628 (1991). Phages can be enriched 20-1,000-fold in a single round of selection. Moreover, the enriched phages can be grown in bacterial culture and subjected to further rounds of selection.

The efficiency of selection depends on many factors, including the kinetics of dissociation during washing, and whether multiple antibody fragments on a single phage can simultaneously engage with antigen. Antibodies with fast dissociation kinetics (and weak binding affinities) can be retained by use of short washes, multivalent phage display and high coating density of antigen in solid phase. The high density not only stabilizes the phage through multivalent interactions, but favors rebinding of phage that has dissociated. The selection of antibodies with slow dissociation kinetics (and good binding affinities) can be promoted by use of long washes and monovalent phage display as described in Bass *et al.*, *Proteins*, **8**: 309-314 (1990) and in WO 92/09690, and a low coating density of antigen as described in Marks *et al.*, *Biotechnol.*, **10**: 779-783 (1992).

It is possible to select between phage antibodies of different affinities, even with affinities that differ slightly, for IFNAR2. However, random mutation of a selected antibody (e.g. as performed in some of the affinity maturation techniques described above) is likely to give rise to many mutants, most binding to antigen, and a few with higher affinity. With limiting IFNAR2, rare high affinity phage could be competed out. To retain all the higher affinity mutants, phages can be incubated with excess biotinylated IFNAR2, but with the biotinylated IFNAR2 at a concentration of lower molarity than the target molar affinity constant for IFNAR2. The high

affinity-binding phages can then be captured by streptavidin-coated paramagnetic beads. Such "equilibrium capture" allows the antibodies to be selected according to their affinities of binding, with sensitivity that permits isolation of mutant clones with as little as two-fold higher affinity from a great excess of phages with lower affinity. Conditions used in washing phages bound to a solid phase can also be manipulated to discriminate on the basis of dissociation kinetics.

5

3. Activity Selection of Anti-IFNAR2 Clones

In one embodiment, the invention provides anti-IFNAR2 antibodies that block the binding between a first type I interferon and IFNAR2 and do not block the binding between a second type I interferon and IFNAR2. Fv clones corresponding to such anti-IFNAR2 antibodies can be selected by (1) isolating anti-IFNAR2 clones from a phage library as described in Section B(I)(2) above, and optionally amplifying the isolated population of phage clones by growing up the population in a suitable bacterial host; (2) selecting a first type I interferon and a second type I interferon against which blocking and non-blocking activity, respectively, is desired; (3) adsorbing the anti-IFNAR2 phage clones to immobilized IFNAR2; (4) using an excess of the first type I interferon to elute the adsorbed clones that recognize IFNAR2-binding determinants which overlap or are shared with the IFNAR2-binding determinants of the first type I interferon; (5) readsorbing the clones isolated from step (4) to immobilized IFNAR2; (6) using an excess of the second type I interferon to elute any undesired clones that recognize IFNAR2-binding determinants which overlap or are shared with the IFNAR2-binding determinants of the second type I interferon; and (7) eluting the clones which remain adsorbed following step (6). The IFNAR2-binding competitions used in this process allow the efficient selection of a phage clone that can block the binding of one selected type I interferon to IFNAR2 and that cannot block the binding of a second selected type I interferon to IFNAR2. Optionally, clones with the desired blocking/non-blocking properties can be further enriched by repeating the selection procedures described herein one or more times.

In another embodiment, the anti-IFNAR2 Fv clone is selected by using a second type I interferon that is IFN- α D in steps (1)-(7) above.

Also provided herein are anti-IFNAR2 Fv clones selected by using a second type I interferon that is IFN- α A in steps (1)-(7) above.

In yet another embodiment, the anti-IFNAR2 Fv clone is selected by using a second type I interferon that is IFN- α B in steps (1)-(7) above.

Additionally provided herein are anti-IFNAR2 Fv clones selected by using a second type I interferon that is IFN- α 1 in steps (1)-(7) above.

Further provided herein are anti-IFNAR2 Fv clones selected by using a second type I interferon that is IFN- β in steps (1)-(7) above.

Also encompassed herein are anti-IFNAR2 Fv clones selected by a first type I interferon that is selected from the group consisting of IFN- α D, IFN- α A, IFN- α G and IFN- α B and using a second type I interferon that is IFN- β in steps (1)-(7) above.

The invention additionally provides anti-IFNAR2 antibodies and Fv clones which block the binding of a more than one selected type I interferon to IFNAR2 and which do not block the binding of another type I interferon to IFNAR2. These Fv clones can be selected by repeating steps (4) and (5) in the above procedure for each type I interferon against which blocking activity is desired, i.e., after eluting clones with an excess of one of the type I interferons against which blocking activity is desired, the eluted clones can be reabsorbed to the immobilized IFNAR2 and then subjected to an excess of another type I interferon against which blocking activity is desired, and the process can be repeated until the remaining clones have been eluted from immobilized IFNAR2 by each species of type I interferon against which blocking activity is desired. In one embodiment, an anti-IFNAR2 Fv clone that blocks the binding of IFN- α D, IFN- α A, IFN- α G and IFN- α B to IFNAR2 and does not block the binding of IFN- β to IFNAR2 is selected by eluting clones adsorbed to immobilized IFNAR2 with each of IFN- α D, IFN- α A, IFN- α G and IFN- α B in consecutive repetitions of steps (4) and (5) followed by subjecting IFNAR2-adsorbed clones to elution with IFN- β in step (6) of the above procedure.

The invention further encompasses anti-IFNAR2 antibodies which bind to specific determinant(s) on IFNAR2. Fv clones corresponding to such anti-IFNAR2 antibodies can be conveniently selected by adsorbing library clones to immobilized IFNAR2 mutants containing Ala substitutions at the specific determinants of interest and recovering library clones which fail to adsorb to immobilized, mutant IFNAR2 (i.e. collected from the column flow-through fractions). Next, the recovered clones are adsorbed to immobilized, wild type IFNAR2, and then the adsorbed clones are recovered, e.g. by elution with excess wild type IFNAR2. The first adsorption step removes clones that bind to IFNAR2 but do not bind to the selected determinant(s), and the second adsorption step removes clones that do not bind to IFNAR2 at all, leaving a population of clones enriched for binding to the selected IFNAR2 determinant(s). The desired clone will exhibit binding activity with wild type IFNAR2 that is greater than the clone's binding activity with the corresponding Ala-substituted IFNAR2 mutant (i.e. a binding level with wild type IFNAR2 that is above the background binding level with mutant IFNAR2). Optionally, the desired clone will exhibit binding activity with the corresponding Ala-substituted IFNAR2 mutant that is less than at or about 90%, or less than at or about 70%, or less than at or about 50%, or less than at or about 30%, or less than at or about 20%, or less than at or about 10%, or at or about 0% of the clone's binding activity with wild type IFNAR2.

Optionally, clones that bind to selected IFNAR2 determinants can be further enriched by repeating the selection procedures described herein one or more times.

In one embodiment, the invention provides anti-IFNAR2 antibodies and Fv clones which bind to a determinant that contains one or more of amino acids 49, 51, 52 and 54 in situ in the sequence of IFNAR2. These Fv clones can be selected by (1) isolating anti-IFNAR2 clones from a phage library as described in Section B(I)(2) above, and optionally amplifying the isolated population of phage clones by growing up the population in a suitable bacterial host; (2) adsorbing anti-IFNAR2 phage clones to an immobilized, mutant IFNAR2 containing Ala substitutions at amino acids 49, 51, 52 and 54 in order to adsorb undesired clones which bind to determinants on IFNAR2 that do not overlap with amino acid positions 49, 51, 52 or 54; (3)

recovering the clones which fail to adsorb to immobilized, mutant IFNAR2 from the flow-through fractions in step (2); (4) readsorbing the recovered clones to immobilized, wild type IFNAR2; and (5) recovering the adsorbed clones by elution with excess IFNAR2.

In another embodiment, the invention provides anti-IFNAR2 antibodies and Fv clones which bind to a determinant that contains one or more of amino acids 68, 71 and 72 in situ in the sequence of IFNAR2. These Fv clones can be selected by (1) isolating anti-IFNAR2 clones from a phage library as described in Section B(I)(2) above, and optionally amplifying the isolated population of phage clones by growing up the population in a suitable bacterial host; (2) adsorbing anti-IFNAR2 phage clones to an immobilized, mutant IFNAR2 containing Ala substitutions at amino acids 68, 71 and 72 in order to adsorb undesired clones which bind to determinants on IFNAR2 that do not overlap with amino acid positions 68, 71 or 72; (3) recovering the clones which fail to adsorb to immobilized, mutant IFNAR2 from the flow-through fractions in step (2); (4) readsorbing the recovered clones to immobilized, wild type IFNAR2; and (5) recovering the adsorbed clones by elution with excess IFNAR2.

In yet another embodiment, the invention provides anti-IFNAR2 antibodies and Fv clones which bind to a determinant that contains one or more of amino acids 74, 77 and 78 in situ in the sequence of IFNAR2. These Fv clones can be selected by (1) isolating anti-IFNAR2 clones from a phage library as described in Section B(I)(2) above, and optionally amplifying the isolated population of phage clones by growing up the population in a suitable bacterial host; (2) adsorbing anti-IFNAR2 phage clones to an immobilized, mutant IFNAR2 containing Ala substitutions at amino acids 74, 77 and 78 in order to adsorb undesired clones which bind to determinants on IFNAR2 that do not overlap with amino acid positions 74, 77 or 78; (3) recovering the clones which fail to adsorb to immobilized, mutant IFNAR2 from the flow-through fractions in step (2); (4) readsorbing the recovered clones to immobilized, wild type IFNAR2; and (5) recovering the adsorbed clones by elution with excess IFNAR2.

In still another embodiment, the invention provides anti-IFNAR2 antibodies and Fv clones which bind to a determinant that contains one or both of amino acids 105 and 109 in situ

in the sequence of IFNAR2. These Fv clones can be selected by (1) isolating anti-IFNAR2 clones from a phage library as described in Section B(I)(2) above, and optionally amplifying the isolated population of phage clones by growing up the population in a suitable bacterial host; (2) adsorbing anti-IFNAR2 phage clones to an immobilized, mutant IFNAR2 containing Ala substitutions at amino acids 105 and 109 in order to adsorb undesired clones which bind to determinants on IFNAR2 that do not overlap with amino acid positions 105 or 109; (3) recovering the clones which fail to adsorb to immobilized, mutant IFNAR2 from the flow-through fractions in step (2); (4) readsorbing the recovered clones to immobilized, wild type IFNAR2; and (5) recovering the adsorbed clones by elution with excess IFNAR2.

In a further embodiment, the invention provides anti-IFNAR2 antibodies and Fv clones which bind to a determinant that contains one or more of amino acids 133, 134, 135 and 139 in situ in the sequence of IFNAR2. These Fv clones can be selected by (1) isolating anti-IFNAR2 clones from a phage library as described in Section B(I)(2) above, and optionally amplifying the isolated population of phage clones by growing up the population in a suitable bacterial host; (2) adsorbing anti-IFNAR2 phage clones to an immobilized, mutant IFNAR2 containing Ala substitutions at amino acids 133, 134, 135 and 139 in order to adsorb undesired clones which bind to determinants on IFNAR2 that do not overlap with amino acid positions 133, 134, 135 or 139; (3) recovering the clones which fail to adsorb to immobilized, mutant IFNAR2 from the flow-through fractions in step (2); (4) readsorbing the recovered clones to immobilized, wild type IFNAR2; and (5) recovering the adsorbed clones by elution with excess IFNAR2.

In an additional embodiment, the invention provides anti-IFNAR2 antibodies and Fv clones which bind to a determinant that contains one or more of amino acids 153, 154 and 156 in situ in the sequence of IFNAR2. These Fv clones can be selected by (1) isolating anti-IFNAR2 clones from a phage library as described in Section B(I)(2) above, and optionally amplifying the isolated population of phage clones by growing up the population in a suitable bacterial host; (2) adsorbing anti-IFNAR2 phage clones to an immobilized, mutant IFNAR2 containing Ala substitutions at amino acids 153, 154 and 156 in order to adsorb undesired clones which bind to

determinants on IFNAR2 that do not overlap with amino acid positions 153, 154 or 156; (3) recovering the clones which fail to adsorb to immobilized, mutant IFNAR2 from the flow-through fractions in step (2); (4) readsorbing the recovered clones to immobilized, wild type IFNAR2; and (5) recovering the adsorbed clones by elution with excess IFNAR2.

Also encompassed herein are anti-IFNAR2 antibodies and Fv clones which bind to a determinant that contains one or more of amino acids 49, 51, 52, and 54 in situ in the sequence of IFNAR2, and contains one or more of amino acids 68, 71, and 72 in situ in the sequence of IFNAR2. These Fv clones can be selected by (1) isolating anti-IFNAR2 clones from a phage library as described in Section B(I)(2) above, and optionally amplifying the isolated population of phage clones by growing up the population in a suitable bacterial host; (2) adsorbing anti-IFNAR2 phage clones to an immobilized, mutant IFNAR2 containing an Ala substitutions at amino acids 49, 51, 52, and 54 in order to adsorb undesired clones which bind to determinants on IFNAR2 that do not overlap with amino acid positions 49, 51, 52 or 54; (3) recovering the clones which fail to adsorb to immobilized, mutant IFNAR2 from the flow-through fractions in step (2); (4) repeating steps (2) and (3) using immobilized, mutant IFNAR2 containing Ala substitutions at amino acid positions 68, 71 and 72 as the adsorbent in order to remove undesired clones which bind to determinants on IFNAR2 that do not overlap with amino acid positions 68, 71 or 72; (5) readsorbing the recovered clones to immobilized, wild type IFNAR2; and (6) recovering the adsorbed clones by elution with excess IFNAR2.

Further provided herein are anti-IFNAR2 antibodies and Fv clones which bind to a determinant that contains one or more of amino acids 49, 51, 52, and 54 in situ in the sequence of IFNAR2, contains one or more of amino acids 68, 71, and 72 in situ in the sequence of IFNAR2, contains one or more of amino acids 74, 77, and 78 in situ in the sequence of IFNAR2, contains one or both of amino acids 105 and 109 in situ in the sequence of IFNAR2, contains one or more of amino acids 133, 134, 135, and 139 in situ in the sequence of IFNAR2, and contains one or more of amino acids 153, 154 and 156 in situ in the sequence of IFNAR2. These Fv clones can be selected by (1) isolating anti-IFNAR2 clones from a phage library as described in Section

B(I)(2) above, and optionally amplifying the isolated population of phage clones by growing up the population in a suitable bacterial host; (2) adsorbing anti-IFNAR2 phage clones to an immobilized, mutant IFNAR2 containing an Ala substitutions at amino acids 49, 51, 52, and 54 in order to adsorb undesired clones which bind to determinants on IFNAR2 that do not overlap with amino acid positions 49, 51, 52 or 54; (3) recovering the clones which fail to adsorb to immobilized, mutant IFNAR2 from the flow-through fractions in step (2); (4) repeating steps (2) and (3) in order to screen the recovered clones for non-adsorption to the corresponding immobilized, Ala-substitution mutant IFNAR2 for each combination of amino acid positions that remains to be tested; (5) readsorbing the recovered clones to immobilized, wild type IFNAR2; and (6) recovering the adsorbed clones by elution with excess IFNAR2.

Additionally provided herein are anti-IFNAR2 antibodies and Fv clones which bind to a determinant that contains one or more of amino acids 133, 134, 135, and 139 in situ in the sequence of IFNAR2, and contains one or more of amino acids 153, 154 and 156 in situ in the sequence of IFNAR2. These Fv clones can be selected by (1) isolating anti-IFNAR2 clones from a phage library as described in Section B(I)(2) above, and optionally amplifying the isolated population of phage clones by growing up the population in a suitable bacterial host; (2) adsorbing anti-IFNAR2 phage clones to an immobilized, mutant IFNAR2 containing an Ala substitutions at amino acids 133, 134, 135, and 139 in order to adsorb undesired clones which bind to determinants on IFNAR2 that do not overlap with amino acid positions 133, 134, 135, or 139; (3) recovering the clones which fail to adsorb to immobilized, mutant IFNAR2 from the flow-through fractions in step (2); (4) repeating steps (2) and (3) in order to screen the recovered clones for non-adsorption to the corresponding immobilized, Ala-substitution mutant IFNAR2 for amino acid positions 153, 154 and 156; (5) readsorbing the recovered clones to immobilized, wild type IFNAR2; and (6) recovering the adsorbed clones by elution with excess IFNAR2.

In other embodiments, the invention provides anti-IFNAR2 antibodies and Fv clones which possess combinations of the differential type I interferon activity inhibiting properties and the IFNAR2 determinant binding properties described herein. Fv clones corresponding to these

embodiments can be selected by using combinations of phage display library screening procedures for selection of clones with unique type I interferon activity inhibition profiles and phage display library screening procedures for selection of clones with unique IFNAR2 determinant binding properties.

For example, the invention provides anti-IFNAR2 antibodies and Fv clones which bind to one or more of amino acid positions 133, 134, 135, and 139 in situ in the sequence of IFNAR2, which bind to one or more of amino acids 153, 154 and 156 in situ in the sequence of IFNAR2, which block the binding of a first type I interferon to IFNAR2, and which do not block the binding of a second type I interferon to IFNAR2. Such Fv clones can be selected by (1) isolating anti-IFNAR2 clones from a phage library as described in Section B(I)(2) above, and optionally amplifying the isolated population of phage clones by growing up the population in a suitable bacterial host; (2) selecting a first type I interferon and a second type I interferon against which blocking and non-blocking activity, respectively, is desired; (3) adsorbing the anti-IFNAR2 phage clones to immobilized IFNAR2; (4) using an excess of the first type I interferon to elute the adsorbed clones that recognize IFNAR2-binding determinants which overlap or are shared with the IFNAR2-binding determinants of the first type I interferon; (5) readsorbing the clones isolated from step (4) to immobilized IFNAR2; (6) using an excess of the second type I interferon to elute any undesired clones that recognize IFNAR2-binding determinants which overlap or are shared with the IFNAR2-binding determinants of the second type I interferon; (7) eluting the clones which remain adsorbed following step (6); (8) adsorbing the eluted clones to an immobilized, mutant IFNAR2 containing an Ala substitution at amino acid positions 133, 134, 135, and 139 in the sequence of IFNAR2 in order to adsorb undesired clones which bind to determinants on IFNAR2 that do not overlap with the amino acid positions 133, 134, 135, or 139; (9) recovering the clones which fail to adsorb to immobilized, mutant IFNAR2 from the flow-through fractions in step (8); and (10) repeating steps (8) and (9) in order to screen the recovered clones for non-adsorption to the corresponding immobilized, Ala-substituted mutant IFNAR2 for amino acid positions 153, 154 and 156.

In a preferred embodiment, the anti-IFNAR2 Fv clone is selected by using a second type I interferon that is IFN- β in steps (1)-(10) of the above procedure.

In another preferred embodiment, the anti-IFNAR2 Fv clone is selected by using a first type I interferon that is selected from the group consisting of IFN- α B, IFN- α G, IFN- α A, and IFN- α D and using a second type I interferon that is IFN- β in steps (1)-(10) of the above.

5 procedure.

In another example, the invention provides anti-IFNAR2 antibodies and Fv clones which bind to one or more of amino acid positions 133, 134, 135, and 139 in situ in the sequence of IFNAR2, which bind to one or more of amino acids 153, 154 and 156 in situ in the sequence of IFNAR2, which block the binding of more than one first type I interferon to IFNAR2, and which do not block the binding of a second type I interferon to IFNAR2. Such Fv clones can be selected by (1) isolating anti-IFNAR2 clones from a phage library as described in Section B(I)(2) above, and optionally amplifying the isolated population of phage clones by growing up the population in a suitable bacterial host; (2) selecting the first type I interferons and the second type I interferon against which blocking and non-blocking activity, respectively, is desired; (3) adsorbing the anti-IFNAR2 phage clones to immobilized IFNAR2; (4) using an excess of one of the first type I interferons to elute the adsorbed clones that recognize IFNAR2-binding determinants which overlap or are shared with the IFNAR2-binding determinants of the first type I interferon; (5) readsorbing the clones isolated from step (4) to immobilized IFNAR2; (6) repeating steps (4) and (5) for each of the remaining type I interferons against which blocking activity is desired; (7) using an excess of the second type I interferon to elute any undesired clones that recognize IFNAR2-binding determinants which overlap or are shared with the IFNAR2-binding determinants of the second type I interferon; (8) eluting the clones which remain adsorbed following step (7); (9) adsorbing the eluted clones to an immobilized, mutant IFNAR2 containing an Ala substitution at amino acid positions 133, 134, 135, and 139 in the sequence of IFNAR2 in order to adsorb undesired clones which bind to determinants on IFNAR2 that do not overlap with the amino acid positions 133, 134, 135, or 139; (10) recovering the

clones which fail to adsorb to the immobilized, mutant IFNAR2 from the flow-through fractions in step (9); and (11) repeating steps (9) and (10) in order to screen the recovered clones for non-adsorption to the corresponding immobilized, Ala-substituted mutant IFNAR2 for amino acid positions 153, 154 and 156.

In a preferred embodiment, the anti-IFNAR2 Fv clone is selected by using a second type I interferon that is IFN- β in steps (1)-(11) of the above procedure.

In another preferred embodiment, the anti-IFNAR2 Fv clone is selected by using all of IFN- α B, IFN- α G, IFN- α A, and IFN- α D as the first type I interferons and using IFN- β as the second type I interferon in steps (1)-(11) of the above procedure.

II. Methods of Making Anti-IFNAR2 Hybridomas

The anti-IFNAR2 antibodies of the invention are preferably monoclonal. Also encompassed within the scope of the invention are Fab, Fab', Fab'-SH and F(ab')₂ fragments of the anti-IFNAR2 antibodies provided herein. These antibody fragments can be created by traditional means, such as enzymatic digestion, or may be generated by recombinant techniques. Such antibody fragments may be chimeric or humanized. These fragments are useful for the diagnostic and therapeutic purposes set forth below.

Monoclonal antibodies are obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Thus, the modifier "monoclonal" indicates the character of the antibody as not being a mixture of discrete antibodies.

The anti-IFNAR2 monoclonal antibodies of the invention can be made using the hybridoma method first described by Kohler *et al.*, *Nature*, **256**:495 (1975), or may be made by recombinant DNA methods (U.S. Patent No. 4,816,567).

In the hybridoma method, a mouse or other appropriate host animal, such as a hamster, is immunized to elicit lymphocytes that produce or are capable of producing antibodies that will

specifically bind to the protein used for immunization. Antibodies to IFNAR2 generally are raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of IFNAR2 and an adjuvant. In one embodiment, animals are immunized with a derivative of IFNAR2 that contains the extracellular domain (ECD) of IFNAR2 fused to the Fc portion of an immunoglobulin heavy chain. In a preferred embodiment, animals are immunized with an IFNAR2-IgG1 fusion protein as described in the Example below. Animals ordinarily are immunized against immunogenic conjugates or derivatives of IFNAR2 with monophosphoryl lipid A (MPL)/trehalose dicrynomycolate (TDM) (Ribi Immunochem. Research, Inc., Hamilton, MT) and the solution is injected intradermally at multiple sites. Two weeks later the animals are boosted. 7 to 14 days later animals are bled and the serum is assayed for anti-IFNAR2 titer. Animals are boosted until titer plateaus.

Alternatively, lymphocytes may be immunized *in vitro*. Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)).

The hybridoma cells thus prepared are seeded and grown in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells. For example, if the parental myeloma cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient cells.

Preferred myeloma cells are those that fuse efficiently, support stable high-level production of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. Among these, preferred myeloma cell lines are murine myeloma lines, such as those derived from MOPC-21 and MPC-11 mouse tumors available from the Salk Institute Cell Distribution Center, San Diego, California USA, and SP-2 or X63-Ag8-653 cells available from the American Type Culture Collection, Rockville, Maryland USA. Human

myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, *J. Immunol.*, **133**:3001 (1984); Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987)).

Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against IFNAR2. Preferably, the binding specificity of monoclonal antibodies produced by hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoadsorbent assay (ELISA).

The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson *et al.*, *Anal. Biochem.*, **107**:220 (1980).

After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may be grown *in vivo* as ascites tumors in an animal.

The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

Anti-IFNAR2 antibodies of the invention possessing the unique properties described in Section B(I) above can be obtained by screening anti-IFNAR2 hybridoma clones for the desired properties by any convenient method. For example, if an anti-IFNAR2 monoclonal antibody that blocks or does not block the binding of certain type I interferons to IFNAR2 is desired, the candidate antibody can be tested in a binding competition assay, such as a competitive binding ELISA, wherein plate wells are coated with IFNAR2, and a solution of antibody in an excess of

the type I interferon of interest is layered onto the coated plates, and bound antibody is detected enzymatically, e.g. contacting the bound antibody with HRP-conjugated anti-Ig antibody or biotinylated anti-Ig antibody and developing the HRP color reaction., e.g. by developing plates with streptavidin-HRP and/or hydrogen peroxide and detecting the HRP color reaction by spectrophotometry at 490 nm with an ELISA plate reader.

5 In another embodiment, the invention provides anti-IFNAR2 monoclonal antibodies that inhibit the anti-viral activity of a first type I interferon and do not inhibit the anti-viral activity of a second type I interferon. Any convenient type I interferon viral infectivity inhibition assay is suitable for use herein. Such assays are well known in the art, and include, for example, type I interferon-induced inhibition of encephalomyocarditis virus (EMC) infectivity in A549 cells as described in *Current Protocols in Immunology*, Coligan, J.E., Kruisbeek, A.M., Margulies, D.H., Shevach, E.M., and Strober, W., eds, Greene Publishing Associates and Wiley-Interscience, (1992), vol. 2, unit 6.9.1. Generally, cells are seeded in attached cell culture plates, grown for 1 day, and then incubated for an additional day in the presence of a predetermined number of units of a selected type I interferon plus various concentrations of the candidate anti-IFNAR2 antibody. Culture supernatants are then removed and cells are challenged with virus and incubated for an additional day. The candidate anti-IFNAR2 antibody that inhibits the anti-viral activity of a selected type I interferon will inhibit more anti-viral activity than the baseline level of anti-viral activity inhibition measured in the presence of an equivalent concentration of control antibody. Optionally, the candidate anti-IFNAR2 antibody that inhibits the anti-viral activity of a selected type I interferon will inhibit at least at or about 30%, or at least at or about 50%, or at least at or about 70%, or at least at or about 80%, or at least at or about 90%, or at least at or about 95%, or at or about 100% of the activity of the type I interferon in the anti-viral activity assay as compared to baseline activity measured in the presence of an equivalent concentration of control antibody. The candidate anti-IFNAR2 antibody that does not inhibit the anti-viral activity of a selected type I interferon will exhibit similar or approximately the same level of anti-viral activity inhibition as a control antibody.

In a preferred embodiment, each type I interferon species used in the viral infectivity assay is titrated to a concentration that provides the same level of inhibition of viral activity as that induced by a preselected number of units of an IFN- α standard. This concentration serves to provide the normalized units of the subject type I interferon species. In order to assess the ability of an anti-IFNAR2 antibody to inhibit the anti-viral activity of various type I interferons, the effective concentration (EC50) of anti-IFNAR2 antibody for inhibiting 50% of a particular type I interferon's anti-viral activity (at the concentration titrated to provide the normalized units of activity) is determined for each type I interferon to be tested. In another preferred embodiment, each type I interferon to be tested is normalized to at least at or about 1 unit/ml, or at or about 1 unit/ml to at or about 1,000 units/ml, or at or about 1 unit/ml to at or about 100 units/ml, of human IFN- α 2. In yet another preferred embodiment, each type I interferon to be tested is normalized to 100 units/ml of the NIH reference standard for recombinant human IFN- α 2 (IFN- α A).

In still another preferred embodiment, the candidate anti-IFNAR2 antibody that does not inhibit the anti-viral activity of a selected type I interferon will exhibit no anti-viral effect at a concentration of at least at or about 1 μ g/ml, or at least at or about 10 μ g/ml, or at least at or about 20 μ g/ml, or at least at or about 30 μ g/ml, or at least at or about 50 μ g/ml, or at least at or about 100 μ g/ml, in the inhibition of EMC infectivity in A549 cells assay described in *Current Protocols in Immunology*, Coligan, J.E., Kruisbeek, A.M., Margulies, D.H., Shevach, E.M., and Strober, W., eds, Greene Publishing Associates and Wiley-Interscience, (1992), vol. 2, unit 6.9.1, wherein each type I interferon is normalized to 100 units/ml of NIH reference standard for recombinant human IFN- α 2 (IFN- α A).

In one aspect, the invention provides anti-IFNAR2 monoclonal antibodies that inhibit the anti-viral activity of a first type I interferon and do not inhibit the anti-viral activity of IFN- α D.

In another aspect, the invention provides anti-IFNAR2 monoclonal antibodies that inhibit the anti-viral activity of a first type I interferon and do not inhibit the anti-viral activity of IFN- α A.

In yet another aspect, the invention provides anti-IFNAR2 monoclonal antibodies that inhibit the anti-viral activity of a first type I interferon and do not inhibit the anti-viral activity of IFN- α B.

In still another aspect, the invention provides anti-IFNAR2 monoclonal antibodies that inhibit the anti-viral activity of a first type I interferon and do not inhibit the anti-viral activity of IFN- α 1.

In a further aspect, the invention provides anti-IFNAR2 monoclonal antibodies that inhibit the anti-viral activity of a first type I interferon and do not inhibit the anti-viral activity of IFN- β .

In an additional aspect, the invention provides anti-IFNAR2 monoclonal antibodies that inhibit the anti-viral activity of a first type I interferon selected from the group consisting of IFN- α D, IFN- α A, IFN- α G and IFN- α B and do not inhibit the anti-viral activity of IFN- β .

If an anti-IFNAR2 monoclonal antibody that binds to a particular IFNAR2 determinant(s) is desired, the candidate antibody can be screened for the presence or absence of differential affinity to wild type IFNAR2 and to mutant IFNAR2 that contains Ala substitution(s) at the determinant(s) of interest as described above. In one aspect, the candidate antibody can be tested for binding to wild type IFNAR2 and mutant IFNAR2 in an immunoprecipitation or immunoadsorption assay. For example, a capture ELISA can be used wherein plates are coated with a given concentration of wild type IFNAR2 or an equal concentration of mutant IFNAR2, the coated plates are contacted with equal concentrations of the candidate antibody, and the bound antibody is detected enzymatically, e.g. contacting the bound antibody with HRP-conjugated anti-Ig antibody and developing the HRP color reaction. The candidate antibody that binds to the particular IFNAR2 determinant(s) of interest will exhibit binding activity with wild type IFNAR2 that is greater than the candidate antibody's binding activity with the corresponding Ala-substituted IFNAR2 mutant (i.e. a binding level with wild type IFNAR2 that is above the background binding level with mutant IFNAR2). Optionally, the candidate antibody that binds to the particular IFNAR2 determinant(s) of interest will exhibit binding activity with

the corresponding Ala-substituted IFNAR2 mutant that is less than about 50%, or less than about 30%, or less than about 20%, or less than about 10%, or less than about 7%, or less than about 6%, or less than about 5%, or less than about 4%, or less than about 3%, or less than about 2%, or less than about 1%, or about 0% of the antibody's binding activity with wild type IFNAR2, e.g. as determined by dividing the HRP color reaction optical density observed for capture ELISA with IFNAR2 mutant adsorbent by the HRP color reaction optical density observed for capture ELISA with wild type IFNAR2 adsorbent.

In other embodiments, the invention provides anti-IFNAR2 antibodies which possess combinations of the type I interferon activity inhibiting and the IFNAR2 determinant binding properties described herein. Anti-IFNAR2 antibodies corresponding to these embodiments can be obtained by using combinations of the type I interferon competitive binding and/or activity inhibition assays described above for selection of antibodies with unique type I interferon inhibiting properties and the immunoprecipitation or immunoadsorption screening procedures described above for selection of antibodies with unique IFNAR2 determinant binding properties.

In one example, the invention provides an anti-IFNAR2 antibody that binds to one or more of amino acid positions 133, 134, 135, and 139 in situ in the sequence of IFNAR2, binds to one or more of amino acids 153, 154 and 156 in situ in the sequence of IFNAR2, inhibits the anti-viral activity of a first type I interferon, and does not inhibit the anti-viral activity of a second type I interferon.

In another example, the invention provides an anti-IFNAR2 antibody that binds to one or more of amino acid positions 133, 134, 135, and 139 in situ in the sequence of IFNAR2, binds to one or more of amino acids 153, 154 and 156 in situ in the sequence of IFNAR2, inhibits the anti-viral activity of a first type I interferon, and does not inhibit the anti-viral activity of IFN- β .

In still another example, the invention provides an anti-IFNAR2 antibody that binds to one or more of amino acid positions 133, 134, 135, and 139 in situ in the sequence of IFNAR2, binds to one or more of amino acids 153, 154 and 156 in situ in the sequence of IFNAR2,

inhibits the anti-viral activity of a first type I interferon selected from the group consisting of IFN- α D, IFN- α A, IFN- α G and IFN- α B, and does not inhibit the anti-viral activity of IFN- β .

In yet another example, the invention provides an anti-IFNAR2 antibody that binds to one or more of amino acid positions 133, 134, 135, and 139 in situ in the sequence of IFNAR2, binds to one or more of amino acids 153, 154 and 156 in situ in the sequence of IFNAR2,
 5 inhibits the anti-viral activity of more than one type I interferon, and does not inhibit the anti-viral activity of another type I interferon.

In a further example, the invention provides an anti-IFNAR2 antibody that binds to one or more of amino acid positions 133, 134, 135, and 139 in situ in the sequence of IFNAR2, binds to one or more of amino acids 153, 154 and 156 in situ in the sequence of IFNAR2, inhibits the anti-viral activity of more than one type I interferon, and does not inhibit the anti-viral activity of IFN- β .

In an additional example, the invention provides an anti-IFNAR2 antibody that binds to one or more of amino acid positions 133, 134, 135, and 139 in situ in the sequence of IFNAR2, binds to one or more of amino acids 153, 154 and 156 in situ in the sequence of IFNAR2,
 10 inhibits the anti-viral activity of IFN- α D, IFN- α A, IFN- α G and IFN- α B, and does not inhibit the anti-viral activity of IFN- β .

In another embodiment, the invention provides the anti-IFNAR2 monoclonal antibody produced by hybridoma cell line 1F3 (ATCC Deposit No.).

In yet another embodiment, the invention provides the anti-IFNAR2 monoclonal antibody
 15 produced by hybridoma cell line 3B7 (ATCC Deposit No.).

In an additional embodiment, the invention provides the anti-IFNAR2 monoclonal antibody produced by hybridoma cell line 1D3 (ATCC Deposit No.).

In still another embodiment, the invention provides anti-IFNAR2 monoclonal antibodies that compete with 1F3 antibody, 3B7 antibody or 1D3 antibody for binding to IFNAR2. Such
 20 competitor antibodies include antibodies that recognize an IFNAR2 epitope that is the same as or overlaps with the IFNAR2 epitope recognized by an antibody selected from the group consisting

of the 1F3, 3B7 and 1D3 antibodies. Such competitor antibodies can be obtained by screening anti-IFNAR2 hybridoma supernatants for binding to immobilized IFNAR2 in competition with labeled 1F3 antibody, 3B7 antibody or 1D3 antibody. A hybridoma supernatant containing competitor antibody will reduce the amount of bound, labeled antibody detected in the subject competition binding mixture as compared to the amount of bound, labeled antibody detected in a control binding mixture containing irrelevant (or no) antibody. Any of the competition binding assays described in Section IV below are suitable for use in the foregoing procedure.

In another aspect, the invention provides an anti-IFNAR2 monoclonal antibody that comprises the complementarity determining regions (CDRs) of the 1F3 antibody. In yet another aspect, the invention provides an anti-IFNAR2 monoclonal antibody that comprises the complementarity determining regions (CDRs) of the 3B7 antibody. In still another aspect, the invention provides an anti-IFNAR2 monoclonal antibody that comprises the complementarity determining regions (CDRs) of the 1D3 antibody. An anti-IFNAR2 monoclonal antibody that comprises the CDRs of 1F3, 3B7 or 1D3 can be constructed by isolating and cloning DNA encoding the variable regions of the 1F3, 3B7 or 1D3 antibody, identifying the CDRs of the 1F3, 3B7 or 1D3 parental antibody, grafting the CDRs onto a template antibody sequence, e.g. a human antibody sequence which is closest to the corresponding murine sequence of the parental antibody, or a consensus sequence of all human antibodies in the particular subgroup of the parental antibody light or heavy chain, and expressing the resulting chimeric light and/or heavy chain variable region sequence(s), with or without accompanying constant region sequence(s), in recombinant host cells as described in Section III below.

III. Methods of Constructing Recombinant Anti-IFNAR2 Antibodies

DNA encoding the hybridoma-derived monoclonal antibodies or phage display Fv clones of the invention is readily isolated and sequenced using conventional procedures (e.g. by using oligonucleotide primers designed to specifically amplify the heavy and light chain coding

regions of interest from hybridoma or phage DNA template). Once isolated, the DNA can be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of the desired monoclonal antibodies in the recombinant host cells. Review articles on recombinant expression in bacteria of antibody-encoding DNA include Skerra *et al.*, *Curr. Opinion in Immunol.*, **5**: 256 (1993) and Pluckthun, *Immunol. Revs.*, **130**: 151 (1992).

DNA encoding the Fv clones of the invention can be combined with known DNA sequences encoding heavy chain and/or light chain constant regions (e.g. the appropriate DNA sequences can be obtained from Kabat *et al.*, supra) to form clones encoding full or partial length heavy and/or light chains. It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species. A Fv clone derived from the variable domain DNA of one animal (such as human) species and then fused to constant region DNA of another animal species to form coding sequence(s) for "hybrid", full length heavy chain and/or light chain is included in the definition of "chimeric" and "hybrid" antibody as used herein. In a preferred embodiment, a Fv clone derived from human variable DNA is fused to human constant region DNA to form coding sequence(s) for all human, full or partial length heavy and/or light chains.

DNA encoding anti-IFNAR2 antibody derived from a hybridoma of the invention can also be modified, for example, by substituting the coding sequence for human heavy- and light-chain constant domains in place of homologous murine sequences derived from the hybridoma clone (e.g. as in the method of Morrison *et al.*, *Proc. Natl. Acad. Sci. USA*, **81**: 6851-6855 (1984)). DNA encoding a hybridoma or Fv clone-derived antibody or fragment can be further modified by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. In this manner, "chimeric" or "hybrid"

antibodies are prepared that have the binding specificity of the Fv clone or hybridoma clone-derived antibodies of the invention.

Typically, such non-immunoglobulin polypeptides are substituted for the constant domains of an antibody of the invention, or they are substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody comprising one antigen-combining site having specificity for IFNAR2 and another antigen-combining site having specificity for a different antigen.

Chimeric or hybrid antibodies also can be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins can be constructed using a disulfide-exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate.

a. Humanized Antibodies

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. It will be appreciated that variable domain sequences obtained from any non-human animal phage display library-derived Fv clone or from any non-human animal hybridoma-derived antibody clone provided as described herein can serve as the "import" variable domain used in the construction of the humanized antibodies of the invention. Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al.*, *Nature*, **321**: 522 (1986); Riechmann *et al.*, *Nature*, **332**: 323 (1988); Verhoeven *et al.*, *Science*, **239**: 1534 (1988)), by substituting non-human animal, e.g. rodent, CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (Cabilly *et al.*, *supra*), wherein substantially less than an intact human variable

domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in non-human animal, e.g. rodent, antibodies.

The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is very important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a non-human animal, e.g. rodent, antibody is screened against the entire library of known human variable-domain sequences. The human sequence which is closest to that of the non-human animal is then accepted as the human framework (FR) for the humanized antibody (Sims *et al.*, *J. Immunol.*, **151**: 2296 (1993); Chothia and Lesk, *J. Mol. Biol.*, **196**: 901 (1987)). Another method uses a particular framework derived from the consensus sequence of all human antibodies of a particular subgroup light or heavy chains. The same framework can be used for several different humanized antibodies (Carter *et al.*, *Proc. Natl. Acad. Sci USA*, **89**: 4285 (1992); Presta *et al.*, *J. Immunol.*, **151**: 2623 (1993)).

It is also important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, i.e., the analysis of residues that influence the ability of the candidate immunoglobulin to bind to its antigen. In this way, FR residues can be selected and combined from the consensus and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is

achieved. In general, the CDR residues are directly and most substantially involved in influencing antigen binding.

b. Human Antibodies

Human anti-IFNAR2 antibodies of the invention can be constructed by combining Fv clone variable domain sequence(s) selected from human-derived phage display libraries with known human constant domain sequences(s) as described above. Alternatively, human monoclonal anti-IFNAR2 antibodies of the invention can be made by the hybridoma method. Human myeloma and mouse-human heteromyeloma cell lines for the production of human monoclonal antibodies have been described, for example, by Kozbor *J. Immunol.*, **133**: 3001 (1984); Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987); and Boerner *et al.*, *J. Immunol.*, **147**: 86 (1991).

It is now possible to produce transgenic animals (e.g. mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region (JH) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. See, e.g., Jakobovits *et al.*, *Proc. Natl. Acad. Sci USA*, **90**: 2551 (1993); Jakobovits *et al.*, *Nature*, **362**: 255 (1993); Bruggermann *et al.*, *Year in Immunol.*, **7**: 33 (1993).

Gene shuffling can also be used to derive human antibodies from non-human, e.g. rodent, antibodies, where the human antibody has similar affinities and specificities to the starting non-human antibody. According to this method, which is also called "epitope imprinting", either the heavy or light chain variable region of a non-human antibody fragment obtained by phage display techniques as described above is replaced with a repertoire of human V domain genes, creating a population of non-human chain/human chain scFv or Fab chimeras. Selection with

antigen results in isolation of a non-human chain/human chain chimeric scFv or Fab wherein the human chain restores the antigen binding site destroyed upon removal of the corresponding non-human chain in the primary phage display clone, i.e. the epitope governs (imprints) the choice of the human chain partner. When the process is repeated in order to replace the remaining non-human chain, a human antibody is obtained (see PCT WO 93/06213 published April 1, 1993).

Unlike traditional humanization of non-human antibodies by CDR grafting, this technique provides completely human antibodies, which have no FR or CDR residues of non-human origin.

c. Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for IFNAR2 and the other is for any other antigen. Exemplary bispecific antibodies may bind to two different epitopes of the IFNAR2 protein. Bispecific antibodies may also be used to localize cytotoxic agents to cells which express IFNAR2. These antibodies possess an IFNAR2-binding arm and an arm which binds the cytotoxic agent (*e.g.* saporin, anti-interferon- α , vinca alkaloid, ricin A chain, methotrexate or radioactive isotope hapten). Bispecific antibodies can be prepared as full length antibodies or antibody fragments (*e.g.* F(ab')₂ bispecific antibodies).

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy chain-light chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, *Nature*, **305**: 537 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures

are disclosed in WO 93/08829 published May 13, 1993, and in Traunecker *et al.*, *EMBO J.*, **10**: 3655 (1991).

According to a different and more preferred approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1), containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance.

In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in WO 94/04690. For further details of generating bispecific antibodies see, for example, Suresh *et al.*, *Methods in Enzymology*, 121:210 (1986).

According to another approach, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the C_H3 domain of an antibody constant domain. In this method, one or more small amino acid side chains from the

interface of the first antibody molecule are replaced with larger side chains (*e.g.* tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (*e.g.* alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

5 Bispecific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (US Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/00373, and EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in US Patent No. 4,676,980, along with a number of cross-linking techniques.

10 Techniques for generating bispecific antibodies from antibody fragments have also been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, *Science*, **229**: 81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the
15 other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

20 Recent progress has facilitated the direct recovery of Fab'-SH fragments from *E. coli*, which can be chemically coupled to form bispecific antibodies. Shalaby *et al.*, *J. Exp. Med.*, **175**: 217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed
25

was able to bind to cells overexpressing the HER2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny *et al.*, *J. Immunol.*, **148(5)**:1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, **90**:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (VH) connected to a light-chain variable domain (VL) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the VH and VL domains of one fragment are forced to pair with the complementary VL and VH domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See Gruber *et al.*, *J. Immunol.*, **152**:5368 (1994).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt *et al. J. Immunol.* **147**: 60 (1991).

IV. Diagnostic Uses of Anti-IFNAR2 Antibodies

The anti-IFNAR2 antibodies of the invention are unique research reagents which provide anti-type I interferon activity templates for use in chemical library screening, wherein the practitioner can use a signal transduction assay as an initial, high volume screen for agents that exhibit an anti-type I interferon activity pattern that is similar to the anti-type I interferon activity pattern of an anti-IFNAR2 antibody of the invention. In this way, candidate agents likely to exhibit a desired type I interferon activity inhibition profile can be obtained with ease, avoiding

prohibitively expensive and logistically impossible numbers of type I interferon induced viral inhibition assays or cell proliferation inhibition assays on large chemical libraries.

In one embodiment, the anti-IFNAR2 antibodies of the invention are used to screen chemical libraries in a Kinase Receptor Activation (KIRA) Assay as described in WO 95/14930 (published 1 June 1995). The KIRA assay is suitable for use herein because ligand binding to the type I interferon receptor complex in situ in on the surface of host cells expressing the receptor induces a rapid increase in the phosphorylation of tyrosine residues in the intracellular domains of both IFNAR1 and IFNAR2 components of the receptor as taught in Platanius and Colamonici, J. Biol. Chem., 269: 17761-17764 (1994). The level of tyrosine phosphorylation can be used as a measure of signal transduction. The effect of an anti-IFNAR2 antibody of the invention on the levels of tyrosine phosphorylation induced by various type I interferons in the KIRA assay can be used as a bench mark activity pattern for comparison to the activity patterns generated by the library compounds in the assay.

The KIRA assay suitable for use herein employs a host cell that expresses the type I interferon receptor (both IFNAR1 and IFNAR2 components of the receptor) and the particular series of type I interferons which define the inhibitor profile of interest. Cells which naturally express the human type I interferon receptor, such as the human Daudi cells and U-266 human myeloma cells described in Colamonici and Domanski, J. Biol. Chem., 268: 10895-10899 (1993), can be used. In addition, cells which are transfected with the IFNAR1 and IFNAR2 components and contain intracellular signaling proteins necessary for type I interferon signal transduction, such as mouse L-929 cells as described in Domanski et al., J. Biol. Chem., 270: 21606-21611 (1995), can be used. In the KIRA assay, the candidate antagonist is incubated with each type I interferon ligand to be tested, and each incubation mixture is contacted with the type I interferon receptor-expressing host cells. The treated cells are lysed, and IFNAR2 protein in the cell lysate is immobilized by capture with solid phase anti-IFNAR2 antibody. Signal transduction is assayed by measuring the amount of tyrosine phosphorylation that exists in the intracellular domain (ICD) of captured IFNAR2 and the amount of tyrosine phosphorylation that

exists in the intracellular domain of any co-captured IFNAR1. Alternatively, cell lysis and immunoprecipitation can be performed under denaturing conditions in order to avoid co-capture of IFNAR1 and permit measurement of IFNAR2 tyrosine phosphorylation alone, e.g. as described in Platanias et al., *J. Biol. Chem.*, 271: 23630-23633 (1996). The level of tyrosine phosphorylation can be accurately measured with labeled anti-phosphotyrosine antibody which identifies phosphorylated tyrosine residues.

In another embodiment, a host cell coexpressing IFNAR1 and a chimeric construct containing IFNAR2 fused at its carboxy terminus to an affinity handle polypeptide is used in the KIRA assay. The chimeric IFNAR2 construct permits capture of the construct from cell lysate by use of a solid phase capture agent (in place of an anti-IFNAR2 antibody) specific for the affinity handle polypeptide. In a preferred embodiment, the affinity handle polypeptide is Herpes simplex virus glycoprotein D (gD) and the capture agent is an anti-gD monoclonal antibody as described in Examples 2 and 3 of WO 95/14930.

In this system, the anti-IFNAR2 antibody of the invention that possesses the type I interferon inhibition activity profile of interest is used as a standard for analysis of the tyrosine phosphorylation patterns generated by the members of the chemical library that is screened. The IFNAR2 ICD tyrosine phosphorylation pattern generated by the anti-IFNAR2 antibody standard is compared to the tyrosine phosphorylation patterns produced in the library screen, and patterns found to match that of the anti-IFNAR2 antibody standard identify candidate agents that are likely to have a type I interferon activity inhibition profile similar to that of the anti-IFNAR2 antibody standard. Accordingly, the anti-IFNAR2 antibody of the invention provides a useful means to quickly and efficiently screen large chemical libraries for compounds likely to exhibit the particular type I interferon activity inhibition profile of the antibody.

In addition, the anti-IFNAR2 antibodies of the invention are useful in diagnostic assays for IFNAR2 expression in specific cells or tissues wherein the antibodies are labeled as described below and/or are immobilized on an insoluble matrix. Anti-IFNAR2 antibodies also are useful for the affinity purification of IFNAR2 from recombinant cell culture or natural sources.

Anti-IFNAR2 antibodies can be used for the detection of IFNAR2 in any one of a number of well known diagnostic assay methods. For example, a biological sample may be assayed for IFNAR2 by obtaining the sample from a desired source, admixing the sample with anti-IFNAR2 antibody to allow the antibody to form antibody/IFNAR2 complex with any IFNAR2 present in the mixture, and detecting any antibody/IFNAR2 complex present in the mixture. The biological sample may be prepared for assay by methods known in the art which are suitable for the particular sample. The methods of admixing the sample with antibodies and the methods of detecting antibody/IFNAR2 complex are chosen according to the type of assay used. Such assays include competitive and sandwich assays, and steric inhibition assays. Competitive and sandwich methods employ a phase-separation step as an integral part of the method while steric inhibition assays are conducted in a single reaction mixture.

Analytical methods for IFNAR2 all use one or more of the following reagents: labeled IFNAR2 analogue, immobilized IFNAR2 analogue, labeled anti-IFNAR2 antibody, immobilized anti-IFNAR2 antibody and steric conjugates. The labeled reagents also are known as "tracers."

The label used is any detectable functionality that does not interfere with the binding of IFNAR2 and anti-IFNAR2 antibody. Numerous labels are known for use in immunoassay, examples including moieties that may be detected directly, such as fluorochrome, chemiluminescent, and radioactive labels, as well as moieties, such as enzymes, that must be reacted or derivatized to be detected. Examples of such labels include the radioisotopes ^{32}P , ^{14}C , ^{125}I , ^3H , and ^{131}I , fluorophores such as rare earth chelates or fluorescein and its derivatives, rhodamine and its derivatives, dansyl, umbelliferone, luciferases, e.g., firefly luciferase and bacterial luciferase (U.S. Pat. No. 4,737,456), luciferin, 2,3-dihydrophthalazinediones, horseradish peroxidase (HRP), alkaline phosphatase, β -galactosidase, glucoamylase, lysozyme, saccharide oxidases, e.g., glucose oxidase, galactose oxidase, and glucose-6-phosphate dehydrogenase, heterocyclic oxidases such as uricase and xanthine oxidase, coupled with an enzyme that employs hydrogen peroxide to oxidize a dye precursor such as HRP,

lactoperoxidase, or microperoxidase, biotin/avidin, spin labels, bacteriophage labels, stable free radicals, and the like.

Conventional methods are available to bind these labels covalently to proteins or polypeptides. For instance, coupling agents such as dialdehydes, carbodiimides, dimaleimides, bis-imidates, bis-diazotized benzidine, and the like may be used to tag the antibodies with the above-described fluorescent, chemiluminescent, and enzyme labels. See, for example, U.S. Pat. Nos. 3,940,475 (fluorimetry) and 3,645,090 (enzymes); Hunter *et al.*, *Nature*, **144**: 945 (1962); David *et al.*, *Biochemistry*, **13**: 1014-1021 (1974); Pain *et al.*, *J. Immunol. Methods*, **40**: 219-230 (1981); and Nygren, *J. Histochem. and Cytochem.*, **30**: 407-412 (1982). Preferred labels herein are enzymes such as horseradish peroxidase and alkaline phosphatase.

The conjugation of such label, including the enzymes, to the antibody is a standard manipulative procedure for one of ordinary skill in immunoassay techniques. See, for example, O'Sullivan *et al.*, "Methods for the Preparation of Enzyme-antibody Conjugates for Use in Enzyme Immunoassay," in *Methods in Enzymology*, ed. J.J. Langone and H. Van Vunakis, Vol. 73 (Academic Press, New York, New York, 1981), pp. 147-166.

Immobilization of reagents is required for certain assay methods. Immobilization entails separating the anti-IFNAR2 antibody from any IFNAR2 that remains free in solution. This conventionally is accomplished by either insolubilizing the anti-IFNAR2 antibody or IFNAR2 analogue before the assay procedure, as by adsorption to a water-insoluble matrix or surface (Bennich *et al.*, U.S. 3,720,760), by covalent coupling (for example, using glutaraldehyde cross-linking), or by insolubilizing the anti-IFNAR2 antibody or IFNAR2 analogue afterward, e.g., by immunoprecipitation.

Other assay methods, known as competitive or sandwich assays, are well established and widely used in the commercial diagnostics industry.

Competitive assays rely on the ability of a tracer IFNAR2 analogue to compete with the test sample IFNAR2 for a limited number of anti-IFNAR2 antibody antigen-binding sites. The anti-IFNAR2 antibody generally is insolubilized before or after the competition and then the

tracer and IFNAR2 bound to the anti-IFNAR2 antibody are separated from the unbound tracer and IFNAR2. This separation is accomplished by decanting (where the binding partner was preinsolubilized) or by centrifuging (where the binding partner was precipitated after the competitive reaction). The amount of test sample IFNAR2 is inversely proportional to the amount of bound tracer as measured by the amount of marker substance. Dose-response curves with known amounts of IFNAR2 are prepared and compared with the test results to quantitatively determine the amount of IFNAR2 present in the test sample. These assays are called ELISA systems when enzymes are used as the detectable markers.

Another species of competitive assay, called a "homogeneous" assay, does not require a phase separation. Here, a conjugate of an enzyme with the IFNAR2 is prepared and used such that when anti-IFNAR2 antibody binds to the IFNAR2 the presence of the anti-IFNAR2 antibody modifies the enzyme activity. In this case, the IFNAR2 or its immunologically active fragments are conjugated with a bifunctional organic bridge to an enzyme such as peroxidase. Conjugates are selected for use with anti-IFNAR2 antibody so that binding of the anti-IFNAR2 antibody inhibits or potentiates the enzyme activity of the label. This method *per se* is widely practiced under the name of EMIT.

Steric conjugates are used in steric hindrance methods for homogeneous assay. These conjugates are synthesized by covalently linking a low-molecular-weight hapten to a small IFNAR2 fragment so that antibody to hapten is substantially unable to bind the conjugate at the same time as anti-IFNAR2 antibody. Under this assay procedure the IFNAR2 present in the test sample will bind anti-IFNAR2 antibody, thereby allowing anti-hapten to bind the conjugate, resulting in a change in the character of the conjugate hapten, e.g., a change in fluorescence when the hapten is a fluorophore.

Sandwich assays particularly are useful for the determination of IFNAR2 or anti-IFNAR2 antibodies. In sequential sandwich assays an immobilized anti-IFNAR2 antibody is used to adsorb test sample IFNAR2, the test sample is removed as by washing, the bound IFNAR2 is used to adsorb a second, labeled anti-IFNAR2 antibody and bound material is then separated

from residual tracer. The amount of bound tracer is directly proportional to test sample IFNAR2. In "simultaneous" sandwich assays the test sample is not separated before adding the labeled anti-IFNAR2. A sequential sandwich assay using an anti-IFNAR2 monoclonal antibody as one antibody and a polyclonal anti-IFNAR2 antibody as the other is useful in testing samples for IFNAR2.

5 The foregoing are merely exemplary diagnostic assays for IFNAR2. Other methods now or hereafter developed that use anti-IFNAR2 antibody for the determination of IFNAR2 are included within the scope hereof, including the bioassays described above.

V. Therapeutic Compositions and Administration of Anti-IFNAR2 Antibodies

10 Therapeutic formulations of the anti-IFNAR2 antibodies of the invention are prepared for storage by mixing antibody having the desired degree of purity with optional physiologically acceptable carriers, excipients, or stabilizers (*Remington: The Science and Practice of Pharmacy*, 19th Edition, Alfonso, R., ed, Mack Publishing Co. (Easton, PA: 1995)), in the form of lyophilized cake or aqueous solutions. Acceptable carriers, excipients or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as Tween, Pluronic or polyethylene glycol (PEG).

20 The anti-IFNAR2 antibody to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution. The anti-IFNAR2 antibody ordinarily will be stored in
25 lyophilized form or in solution.

Therapeutic anti-IFNAR2 antibody compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

The route of anti-IFNAR2 antibody administration is in accord with known methods, e.g. injection or infusion by intravenous, intraperitoneal, intracerebral, subcutaneous, intramuscular, intraocular, intraarterial, intracerebrospinal, or intralesional routes, or by sustained release systems as noted below. Preferably the antibody is given systemically.

Suitable examples of sustained-release preparations include semipermeable polymer matrices in the form of shaped articles, e.g. films, or microcapsules. Sustained release matrices include polyesters, hydrogels, polylactides (U.S. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman *et al.*, *Biopolymers*, **22**: 547-556 (1983)), poly (2-hydroxyethyl-methacrylate) (Langer *et al.*, *J. Biomed. Mater. Res.*, **15**: 167-277 (1981) and Langer, *Chem. Tech.*, **12**: 98-105 (1982)), ethylene vinyl acetate (Langer *et al.*, *supra*) or poly-D-(-)-3-hydroxybutyric acid (EP 133,988). Sustained-release anti-IFNAR2 antibody compositions also include liposomally entrapped antibody. Liposomes containing antibody are prepared by methods known *per se*: DE 3,218,121; Epstein *et al.*, *Proc. Natl. Acad. Sci. USA*, **82**: 3688-3692 (1985); Hwang *et al.*, *Proc. Natl. Acad. Sci. USA*, **77**: 4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese patent application 83-118008; U.S. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. % cholesterol, the selected proportion being adjusted for the optimal antibody therapy.

Anti-IFNAR2 antibody can also be administered by inhalation. Commercially available nebulizers for liquid formulations, including jet nebulizers and ultrasonic nebulizers are useful for administration. Liquid formulations can be directly nebulized and lyophilized powder can be nebulized after reconstitution. Alternatively, anti-IFNAR2 antibody can be aerosolized using a fluorocarbon formulation and a metered dose inhaler, or inhaled as a lyophilized and milled powder.

An "effective amount" of anti-IFNAR2 antibody to be employed therapeutically will depend, for example, upon the therapeutic objectives, the route of administration, the type of anti-IFNAR2 antibody employed, and the condition of the patient. Accordingly, it will be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. Typically, the clinician will administer the anti-IFNAR2 antibody until a dosage is reached that achieves the desired effect. The progress of this therapy is easily monitored by conventional assays.

The patients to be treated with the anti-IFNAR2 antibody of the invention include preclinical patients or those with recent onset of immune-mediated disorders, and particularly autoimmune disorders. Patients are candidates for therapy in accord with this invention until such point as no healthy tissue remains to be protected from immune-mediated destruction. For example, a patient suffering from insulin-dependent diabetes mellitus (IDDM) can benefit from therapy with an anti-IFNAR2 antibody of the invention until the patient's pancreatic islet cells are no longer viable. It is desirable to administer an anti-IFNAR2 antibody as early as possible in the development of the immune-mediated or autoimmune disorder, and to continue treatment for as long as is necessary for the protection of healthy tissue from destruction by the patient's immune system. For example, the IDDM patient is treated until insulin monitoring demonstrates adequate islet response and other indicia of islet necrosis diminish (e.g. reduction in anti-islet antibody titers), after which the patient can be withdrawn from anti-IFNAR2 antibody treatment for a trial period during which insulin response and the level of anti-islet antibodies are monitored for relapse.

In the treatment and prevention of an immune-mediated or autoimmune disorder by an anti-IFNAR2 antibody, the antibody composition will be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the antibody, the particular type of antibody, the method of administration, the scheduling of administration, and

other factors known to medical practitioners. The "therapeutically effective amount" of antibody to be administered will be governed by such considerations, and is the minimum amount necessary to prevent, ameliorate, or treat the disorder, including treating chronic autoimmune conditions and immunosuppression maintenance in transplant recipients. Such amount is preferably below the amount that is toxic to the host or renders the host significantly more susceptible to infections.

As a general proposition, the initial pharmaceutically effective amount of the antibody administered parenterally will be in the range of about 0.1 to 50 mg/kg of patient body weight per day, with the typical initial range of antibody used being 0.3 to 20 mg/kg/day, more preferably 0.3 to 15 mg/kg/day. The desired dosage can be delivered by a single bolus administration, by multiple bolus administrations, or by continuous infusion administration of antibody, depending on the pattern of pharmacokinetic decay that the practitioner wishes to achieve.

As noted above, however, these suggested amounts of antibody are subject to a great deal of therapeutic discretion. The key factor in selecting an appropriate dose and scheduling is the result obtained, as indicated above.

The antibody need not be, but is optionally formulated with one or more agents currently used to prevent or treat the immune-mediated or autoimmune disorder in question. For example, in rheumatoid arthritis, the antibody may be given in conjunction with a glucocorticosteroid. The effective amount of such other agents depends on the amount of anti-IFNAR2 antibody present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as used hereinbefore or about from 1 to 99% of the heretofore employed dosages.

Further details of the invention can be found in the following example, which further defines the scope of the invention. All references cited throughout the specification, and the references cited therein, are hereby expressly incorporated by reference in their entirety.

EXAMPLE

MATERIALS AND METHODS

Preparation of soluble IFNAR2-IgG.

A cDNA encoding the human immunoglobulin fusion proteins (immunoadhesins) based on the ECD of the hIFNAR2 (pRK5 hIFNAR2-IgG clone) was generated using methods similar to those described by Haak-Frendscho *et al.*, *Immunology* **79**: 594-599 (1993) for the construction of a murine IFN- γ receptor immunoadhesin. Briefly, the plasmid pRKCD4₂Fc₁ was constructed as described in Example 4 of WO 89/02922 (PCT/US88/03414 published April 6, 1989). The cDNA coding sequence for the first 216 residues of the mature hIFNAR2 ECD shown in Fig. 5 was obtained from the published sequence (Novick *et al.*, *Cell*, **77**: 391-400 (1994)). The CD4 coding sequence in the pRKCD4₂Fc₁ was replaced with the hIFNAR2 ECD encoding cDNA to form the pRK5hIFNAR2-IgG clone. The nucleic acid sequence (SEQ ID NO. 25) and amino acid sequence (SEQ ID NO. 26) for the hIFNAR2 ECD-IgG encoding insert of the clone are shown in Fig. 5. hIFNAR2-IgG was expressed in human embryonic kidney 293 cells by transient transfection using a calcium phosphate precipitation technique. The immunoadhesin was purified from serum-free cell culture supernatants in a single step by affinity chromatography on a protein A-sepharose column as described in Haak-Frendscho *et al.* (1993), *supra*. Bound hIFNAR2-IgG was eluted with 0.1 M citrate buffer, pH 3.0, containing 20% (w/v) glycerol. The hIFNAR2-IgG purified was over 95% pure, as judged by SDS-PAGE.

Production of hIFN- α subtypes.

Standard cloning procedures described in Maniatis *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989) were used to construct plasmids that direct the translocation of the various species of hIFN- α into the periplasmic space of *E. coli*. PCR reactions were performed on cDNA clones of the various subspecies of hIFN- α disclosed in Goeddel *et al.*, *Nature* **290**: 20-26 (1981) with Nsil and StyI restriction sites added to the primers. These PCR products were then subcloned into the

corresponding sites of the expression vector pB0720 described in Cunningham *et al.*, *Science* **243**:1330-1336 (1989). The resulting plasmids placed production of the hIFN- α subtypes under control of the *E. coli* *phoA* promoter and the heat-stable enterotoxin II signal peptide as described in Chang *et al.*, *Gene* **55**: 189-196 (1987). The correct DNA sequence of each gene was confirmed using the United States Biochemical Sequenase Kit version 2.0. Each plasmid was transformed into the *E. coli* strain 27C7 (ATCC # 55244) and grown in 10 liter fermentors as described in Carter *et al.*, *Bio/Technology* **10**: 163-167 (1992). Human hIFNs were purified from *E. coli* paste containing each IFN- α by affinity chromatography. Bacterial cells were lysed, and the lysate was centrifuged at 10,000 x g to remove debris. The supernatant was applied to an immunoaffinity column containing a mouse anti-hIFN- α B antibody (LI-1) that was obtained as described in Staehelin *et al.*, *Proc. Natl. Acad. Sci.* **78**:1848-1852 (1981). LI-1 was immobilized on controlled pore glass by a modification of the method of Roy *et al.*, *Journal of Chromatography*, **303**: 225-228 (1984). The bound interferon was eluted from the column with 0.1 M citrate, pH 3.0, containing 20% (w/v) glycerol. The purified IFN was analyzed by SDS-PAGE and immunoblotting, and was assayed for bioactivity by the hIFN-induced anti-viral assay as described herein. hIFN β was obtained from Sigma (St. Louis, Mo.).

Human IFN- α 2/1 hybrid molecule (IFN- α 2₁₋₆₂/ α 164-166) was obtained as described in Rehberg *et al.*, *J. Biol. Chem.*, **257**: 11497-11502 (1992) or Horisberger and Marco, *Pharmac. Ther.*, **66**: 507-534 (1995).

Generation of mAbs to hIFNAR2.

Balb/c mice were immunized into each hind foot pad 11 times at two week intervals, with 2.5 μ g of hIFNAR2-IgG fusion protein resuspended in MPL-TDM (Ribi Immunochem. Research Inc., Hamilton, MT). Three days after the final boost, popliteal lymph node cells were fused with murine myeloma cells, P3X63AgU.1 (ATCC CRL1597), using 35% polyethylene glycol.

Hybridomas were selected in HAT medium. Ten days after the fusion, hybridoma culture

supernatants were first screened for mAbs binding to the hIFNAR2-IgG fusion protein but not to CD4-IgG in a capture ELISA. The selected culture supernatants further tested for their ability to block the ligand-receptor binding in a capture ELISA as described below and for their ability to recognize cell membrane receptors on U266 cells by flow cytometric analysis as described in Chuntharapai *et al.*, *J. Immunol.*, **152**:1783-1789 (1994). After cloning the selected final
5 hybridomas twice, their antigen specificity as well as blocking activities were confirmed in the ligand-receptor binding assay, IGSF-3 complex assay and anti-viral assay as described below.

Epitope mapping using a competitive binding ELISA.

To determine whether the mAbs recognized the same or different epitopes, a competitive binding ELISA was performed as described in Kim *et al.*, *J. Immunol. Method* **156**: 9-17 (1992), using biotinylated mAbs (Bio-mAb). mAbs were biotinylated using N-hydroxyl succinimide as described in *Antibodies (A Laboratory Manual)*, Harlow, E. and Lane, D., eds, Cold Spring Harbor (1988), p. 341. Microtiter wells were coated with 50 µl of Goat anti-hIgG-Fc and kept overnight at 4°C, blocked with assay buffer for 1 hour, and incubated with 25 µl/well of IFNAR2-IgG (1 µg/ml) for 1 hour at room temperature. After washing microtiter wells, a mixture of a predetermined optimal concentration of Bio-mAb and a thousand-fold excess of unlabeled mAb was added into each well. Following 1 hour incubation at room temperature, plates were washed and the amount of Bio-mAb was detected by the addition of HRP-streptavidin. After washing the microtiter wells, the bound enzyme was detected by the addition
20 of the substrate, and the plates were read at 490 nm with an ELISA plate reader.

Determination of the affinities of mAbs

The equilibrium dissociation and association constant rates of anti-IFNAR2 mAbs were
25 determined using KinExATM automated immunoassay system (Sapidyne Instruments, Inc. Boise, ID) modified as described in Blake *et al.*, *J. Biol. Chem.*, **271**: 27677 (1996). Briefly, 1.0 ml of

Anti-Human IgG Agarose beads (56 μm , Sigma. St.Louis, MO.) were coated with 20 μg of IFNAR2-IgG in PBS by gentle mixing at room temperature (RT) for 1 hour. After washing with PBS, non specific binding sites were blocked by incubating with 10 % human serum in PBS for 1 hour at RT. A bead pack (approximately 4 mm high) was created in the observation flow cell of the KinExATM instrument. Briefly, the blocked beads were diluted into 30 ml of assay buffer (0.01 % BSA/PBS). The diluted beads (550 μl) were drawn through the flow cell with a 20 micron screen and then washed with 1 ml of running buffer (0.01% BSA + 0.05% Tween 20 in PBS). The beads were then disrupted gently with a brief backflush of running buffer, and allowed to set for 20 seconds in order to create a uniform and reproducible bead pack.

For equilibrium measurements, mAbs (5 ng/ml-31 pM-in 0.01 % BSA/PBS)) were mixed with a serial dilutions of IFNAR2-IgG (concentrations from 2.5 nM to 5.0 pM) and incubated at RT for 2 minutes. Once equilibrium was reached, 4.5 ml of this mixture was drawn through the beads, followed by 250 μl of running buffer to wash out the unbound mAb. The primary mAbs bound to beads were detected by 1.5 ml of phycoerythrin (PE) labeled goat anti-mouse IgG drawn through the bead pack. Unbound labeled material was removed by drawing 4.5 ml of 0.5 M NaCl through the bead pack over a 3 minute period. The equilibrium constant was calculated by Scatchard analysis as described in Munson et al., Anal. Biochem., **160**: 1085 (1980).

Electrophoretic Mobility Shift Assay (EMSA)

Briefly, 5 ng of α -IFNs plus various concentrations of anti-hIFNAR2 mAbs were incubated with 5×10^5 Hela cells in 200 μl of DMEM for 30 minutes at 37° C. Cells were washed in PBS and resuspended in 125 μl buffer A (10 mM HEPES, pH 7.9, 10 mM KCL, 0.1 mM EDTA, 1 mM DTT, 1 mM phenylmethylsulfonyl floride, 10 $\mu\text{g}/\text{ml}$ leupeptin, 10 $\mu\text{g}/\text{ml}$ aprotinin) as described in Kurabayashi et al., Mol. Cell Biol., **15**: 6386 (1995). After a 15 minute incubation on ice, cells were lysed by the addition of 0.025% NP40. The nuclear pellet was obtained by centrifugation and was resuspended in 50 μl buffer B (20 mM HEPES, pH 7.9, 400 mM NaCL, 0.1 mM EDTA, 1 mM DTT, 1 mM phenylmethylsulfonyl floride, 10 $\mu\text{g}/\text{ml}$

leupeptin, 10 µg/ml aprotinin) and kept on ice for 30 min. The nuclear fraction was cleared by centrifugation and the supernatant stored at -70°C until use. Double-stranded probes were prepared from single-stranded oligonucleotides (ISG15 top : 5'-GATCGGGAAAGGGAAACCGAAACTGAAGCC-3' (SEQ ID NO. 1)), ISG15 bottom: 5'-GATCGGCTTCAGTTTCGGTTTCCCTTTCCC-3' (SEQ ID NO. 2)) utilizing a DNA
 5 polymerase I Klenow filling reaction with ³²P- dATP (3,000 Ci/mM, Amersham). Labeled oligonucleotides were purified from unincorporated radioactive nucleotides using BIO-Spin 30 columns (Bio-Rad). Binding reactions containing 5 µl nuclear extract, 25,000 cpm of labeled probe and 2 µg of non-specific competitor poly (dI-dC)-poly (dI-dC) in 15 µl binding buffer (10 mM Tris-HCL, pH 7.5, 50 mM NaCl, 1 mM EDTA, 1 mM DTT, 1 mM phenylmethylsulfonyl
 10 floride and 15% glycerol) were incubated at RT for 30 minutes. DNA-protein complexes were resolved in 6% non-denaturing polyacrylamide gels and analyzed by autoradiograph. The specificity of the assay was determined by the addition of 350 ng of unlabeled ISG15 probe in separate reaction mixtures. Formation of an ISGF3 specific complex was confirmed by a super
 15 shift assay with anti-STAT1 antibody.

Anti-viral assay

The assay was done as described in *Current Protocols in Immunol.*, Coligan, J.E., Kruisbeek, A.M., Margulies, D.H., Shevach, E.M., and Strober, W., eds, Greene Publishing Associates and Wiley-Interscience (1992), Vol. 2, Unit 6.9.1, using the human lung carcinoma
 20 cell line A549 challenged with encephalomyocarditis virus (EMC). Serial dilutions of mAbs were incubated with various units of type I IFNs in 50 µl DMEM at 37°C. These mixtures were then incubated with A549 cells (5x10⁵ cells/100 µl DMEM 4% FCS) for another 24 hours. Culture supernatants were then removed and challenged with 2x10⁵ pfu of encephalomyocarditis (EMC) virus in 100 µl DMEM with 2% FCS for an additional 24 hours. At the end of the
 25 incubation, cell viability was determined by visual microscopic examination. Cells were then incubated with seven 1:2 dilutions of mixtures containing anti-IFNAR2 mAbs plus type 1 hIFN

for 24 hours. Each dilution was tested in duplicate. Cells were then washed and challenged with encephalomyocarditis (EMC) virus for another 24 hours. At the end of the experiment, the remaining viable cells were detected by crystal violet staining. The neutralizing antibody titer (EC50) was defined as the concentration of antibody which neutralizes 50% of the anti-viral cytopathic effect induced by 100 units/ml of type I IFNs. The units of type I IFNs used in this study were determined using NIH reference human IFN- α 2 as a standard.

Generation of various hIFNAR2-IgG mutants

A cDNAs encoding residues 1-216 of the extracellular domain of IFNAR2 was constructed and expressed as an immunoadhesin. Single alanine substitution mutants were generated according to the method of Kunkel *et al.*, *Methods Enzymol.* **154**: 367-414 (1987), and Hebert *et al.*, *J. Biol. Chem.*, **268**: 18549-18553 (1993). The plasmid DNA was isolated using an RPM Kit (BIO 101 Inc., La Jolla, CA) and was sequenced by the Sanger method using an ABI 373A DNA sequencer to verify the mutation. Mutant receptor-IgGs were expressed transiently in human 293 cells as described above. Transfected 293 cells were grown overnight in F-12:DMEM (50:50) containing 10% FCS, 2 mM glutamine, 100 μ g/ml of penicillin, 100 μ g/ml of streptomycin, 10 μ g/ml of glycine, 15 μ g/ml of hypoxanthine, and 5 μ g/ml of thymidine, and then were placed in serum-free media. Three days later, culture supernatants were collected and used in a capture ELISA to determine the mAb binding sites.

Capture ELISA.

Microtiter plates (Maxisorb; Nunc, Kamstrup, Denmark) were coated with 50 μ l/well of 2 μ g/ml of goat antibodies specific to the Fc portion of human IgG (Goat anti-hIgG-Fc, Cappel), in PBS, overnight at 4°C and blocked with 2% BSA for 1 hour at room temperature. After washing the plate, 50 μ l/well of 2 μ g/ml of IFNAR2-IgG (or IFNAR2-IgG mutant) were added to each well for 1 hour. The remaining anti-Fc binding sites were blocked with PBS containing 3%

human serum and 10 µg/ml of CD4-IgG for 1 hour. Plates were then incubated with 50 µl/well of 2 µg/ml of anti-IFNAR2 mAbs (or hybridoma culture supernatants) for 1 hour. Plates were then incubated with 50 µl/well of HRP-Goat anti-mouse IgG. The bound enzyme was detected by the addition of the TMB (3,3',5,5'-tetramethylbenzidin) substrate, the reaction was stopped by the addition of the stop solution (Kirkegaard & Perry Lab, Gathersburg, MD) and the plates were
5 read at 450 nM with an ELISA plate reader. Between each step, plates were washed three times in wash buffer (PBS containing 0.05% Tween 20).

During the IFNAR2-IgG mutant analysis, the concentrations of immunoadhesin molecules in 293 transfected culture supernatants were determined using CD4-IgG as a standard and were adjusted to be equal to the lowest concentration (approximately 0.5 nm) of immunoadhesin molecules. The degree of mAb binding to these mutants was then compared to the wild type of the same concentration.

RESULTS

mAbs to IFNAR2 binding different epitopes show differential blocking activities.

mAbs 1D3, 1F3 and 3B7, which recognize different epitopes on IFNAR2 by competitive binding ELISA using the molar ratio of biotinylated mAb to unlabeled mAb of 1:100 (Table 1 below), were developed to characterize the structure-function of IFNAR2. All of these mAbs belong to the IgG2a isotype and recognize membrane IFNAR2 on human myeloma U266 cells by FACS.

Table 1
Summary of the general characteristics of mAbs to IFNAR2

mAb	Isotype ^a	FACS ^b	Epitope ^c	Western ^e blot	Imm. ppt ^d	Affinity ^e Kd ⁻¹ (pM)
1D3	IgG2a	+++	A	+	ND	242
1F3	IgG2a	+++	B	-	+	5
3B7	IgG2a	+++	C	-	+	<1

a. The isotype of mAbs was determined using isotype specific goat anti-mouse Ig.

b. All these mAbs stained U266 human myeloma cells expressing IFNAR by FACS.

c. MAbs were shown to recognize different epitopes by competitive binding ELISA.

d. The immunoblotting was performed using IFNAR2 reduced with dithiothreitol (DTT).

e. U266 cells were biotinylated using NHS-LC-biotin and lysed with 1% NP-40. Biotinylated IFNAR2 were then precipitated by mAbs bound to protein-G-4B sepharose and separated on a 7.5% SDS-PAGE gel. Biotinylated IFNAR2 transferred onto nitrocellulose paper was then detected by HRP-streptavidin as described.

f. The affinity of mAbs were determined by Scatchard analysis using KINEX system.

Only mAb 1D3 recognizes the reduced IFNAR2 in the immunoblot, indicating that mAb 1D3 recognizes a linear epitope while others recognize conformational epitopes. The affinities (Kd⁻¹) of mAb 3B7, 1F3 and 1D3 were less than 1 pM, 5 pM and 242 pM, respectively, demonstrating that these are relatively high affinity mAbs

The neutralizing abilities of these mAbs were determined in the receptor-ligand binding ELISA (Fig 1). At a concentration of 0.6 nM (0.1 µg/ml), mAbs 1F3 and 3B7 were able to block greater than 90% of the biotinylated IFN α -2/1 binding to IFNAR2-IgG while even at a concentration of 6 nM (1 µg/ml), mAb 1D3 showed no significant blocking activity. From these ELISA results, it was determined that mAbs 1F3 and 3B7 are blocking mAbs and that mAb 1D3 may not be a blocking mAb. To further determine blocking activities of these mAbs, mAbs were tested in the IGSF complex formation EMSA and in the anti-viral assay. Fig. 2 depicts the results obtained in IGSF complex formation induced by IFN- α 2 (IFN- α A) using L929 cells expressing hIFNAR2. At a concentration of 1 µg/ml, mAbs 3B7 and 1F3, but not mAb 1D3, completely blocked the IGSF complex formation induced by human type 1 IFN- α 2. At a

concentration of 10 µg/ml, all three mAbs blocked the IGSF complex formation. These results demonstrated that all three mAbs are blocking mAbs but mAb 1D3 is a weak blocking mAb. Blocking activities of these mAbs on the anti-viral activities of type 1 IFNs (IFN α -1(D), -2(A), -5(G), -8(B), -2/1 and IFN- β) are summarized in Table 2 below.

Table 2.
Effects of anti-hIFNAR2 mAbs on the anti-viral effects of type 1 IFNs

mAb	EC50 of mAb (µg/ml)					
	IFN α 2/1	IFN α 1	IFN α 2	IFN α 5	IFN α 8	IFN β
1D3	20	10	20	10	20	NB
1F3	3	2	3	1	3	2
3B7	0.6	0.1	1	0.1	0.3	0.3

The neutralizing antibody titer (EC50) was defined as the concentration of antibody which neutralizes 50% of the anti-viral cytopathic effects induced by 100 units/ml of type 1 IFNs on A549 cells. Experiment were done using serial dilutions of mAbs in the range of 0.1-30 µg/ml in duplicate. mAbs exhibiting no blocking effect at a concentration of 30 µg/ml in the assay were designated as nonblocking mAbs (NB).

mAbs 3B7 and 1F3 blocked the activities of all type 1 IFNs tested. The mAb 3B7 exhibited a blocking activity of 50% reduction in interferon-induced anti-viral cytopathic effects (EC50) at a concentration of less than 1 µg/ml on all type IFNs, and the EC50 concentration of mAb 1F3 was 5-20 fold higher than the EC50 concentration of mAb 3B7. Although mAb 1D3 was able to block the activity of all IFN α species except IFN β , a much higher concentration of 1D3 antibody was required for inhibition of the IFN- α activities compared to mAbs 3B7 and 1F3.

Determination of residues on IFNAR2 which are important for the binding of IFN α -2/1 and neutralizing mAbs. In this study, hIFN- α 2/1 hybrid molecule (IFN- α 2₁₋₆₂/ α 1₆₄₋₁₆₆) was selected for testing due to its availability in a large quantity, its potent anti-viral activities in various animal species and its wide clinical use. The IFN- α 2/1 hybrid molecule was generated by

recombining IFN- α 2(IFN α A) and IFN- α 1(IFN- α D), which are the most common IFN- α species produced in virus-infected leukocytes and which share a high degree of structural homology. The recombinant IFN-2/1 hybrid retains potent anti-viral activities in human as well as in mouse, confirming the presence of functional IFN receptor binding sites.

To determine areas of IFNAR2 which are important for the ligand binding, multiple and single alanine substitution mutants were generated as shown in Tables 3 and 4 below.

5

Table 3.

Binding of IFN- α 2/1 and anti-IFNAR2 mAbs
to multiple alanine IFNAR2 mutants

residues	Alanine substitution	% Wild IFNAR2 binding			
		IFN α -2/1	1D3	1F3	3B7
7-11	DYTDE (SEQ ID NO. 3) / AYTAA (SEQ ID NO. 4)	73 \pm 10	105 \pm 13	96 \pm 1	101 \pm 13
29-33	ELKNH (SEQ ID NO. 5) / ALANA (SEQ ID NO. 6)	87 \pm 22	80 \pm 1	87 \pm 6	82 \pm 7
49-55	KPEDLK (SEQ ID NO. 7) / APAALA (SEQ ID NO. 8)	18 \pm 2	39 \pm 1	6 \pm 0	4 \pm 0
68-72	DLTDE (SEQ ID NO. 9) / ALTAA (SEQ ID NO. 10)	16 \pm 1	38 \pm 2	5 \pm 0	3 \pm 0
74-78	RSTHE (SEQ ID NO. 11) / ASTAA (SEQ ID NO. 12)	16 \pm 1	95 \pm 1	16 \pm 1	89 \pm 2
105-109	DMSFE (SEQ ID NO. 13) / AMSFA (SEQ ID NO. 14)	19 \pm 2	65 \pm 1	8 \pm 1	40 \pm 1
133-139	EEELQFD (SEQ ID NO. 15) / AAALQFA (SEQ ID NO. 16)	16 \pm 1	5 \pm 1	9 \pm 0	35 \pm 1
145-149	EEQSE (SEQ ID NO. 17) / AAQSA (SEQ ID NO. 18)	No expression			
153-157	KKHKP (SEQ ID NO. 19) / AAHAP (SEQ ID NO. 20)	19 \pm 1	14 \pm 1	14 \pm 0	35 \pm 1
159-163	EIKGN (SEQ ID NO. 21) / AIAGN (SEQ ID NO. 22)	92 \pm 5	89 \pm 7	96 \pm 5	93 \pm 3
172-173	DK/ AA	73 \pm 7	73 \pm 5	91 \pm 7	80 \pm 3
187-192	EHSDEA (SEQ ID NO. 23) / AASAAQ (SEQ ID NO. 24)	82 \pm 3	82 \pm 10	83 \pm 9	66 \pm 10

Wild and mutant IFNAR2 adhesin molecules (0.5 nM) were captured with the goat anti-human IgG Fc reagent precoated onto ELISA plate wells. Biotinylated IFN α -A (50 nM) or mAbs (5 nM) were allowed to interact for 1 hour. After washing, the amounts of the ligand or mAbs bound were determined by the addition of HRP-streptavidin and HRP-sheep anti-mouse IgG, respectively.

Table 4

Binding of IFN- α 2/1 and anti-IFNAR2 mAbs to single alanine IFNAR2 mutants

mutants	% Wild IFNAR2 binding				
	IFN- α 2/1	1D3	1F3	3B7	Polyclonal
K49A	68 \pm 10	101 \pm 1	104 \pm 6	99 \pm 1	98 \pm 13
E51A	95 \pm 9	103 \pm 4	101 \pm 4	67 \pm 0	92 \pm 3
D52A	97 \pm 11	101 \pm 1	107 \pm 6	83 \pm 2	94 \pm 15
K54A	53 \pm 1	87 \pm 1	83 \pm 2	87 \pm 12	91 \pm 13
K57A	113	110	120	91	120
D68A	15 \pm 2	41 \pm 1	8 \pm 1	8 \pm 0	38 \pm 1
D71A	113 \pm 26	100 \pm 2	91 \pm 3	110 \pm 13	103 \pm 12
E72A	129 \pm 29	91 \pm 7	106 \pm 19	90 \pm 13	94 \pm 2
R74A	122 \pm 28	92 \pm 2	103 \pm 12	93 \pm 16	93 \pm 13
H77A	83 \pm 19	150 \pm 4	12 \pm 137	132 \pm 16	150 \pm 12
E78A	36 \pm 3	92 \pm 3	54 \pm 4	94 \pm 2	89 \pm 14
W101A	22 \pm 3	99 \pm 4	90 \pm 2	98 \pm 2	100 \pm 13
I104A	51 \pm 5	104 \pm 1	94 \pm 6	104 \pm 1	99 \pm 13
D105A	53 \pm 11	89 \pm 3	62 \pm 8	87 \pm 2	97 \pm 14
E109A	70 \pm 9	106	121	111	100 \pm 17
E133A	78	92 \pm 3	96 \pm 12	82	80 \pm 12
E134A	86 \pm 3	87 \pm 1	82 \pm 3	88 \pm 12	92 \pm 3
E135A	72 \pm 2	88 \pm 1	74 \pm 1	80 \pm 12	90 \pm 1
Q137A	43 \pm 4	63 \pm 1	55 \pm 4	61 \pm 14	82 \pm 1
D139A	62	87 \pm	100 \pm 4	100	89 \pm 14
E145A	78 \pm 9	84 \pm 2	82 \pm 12	93 \pm 12	91 \pm 11
E146A	87 \pm 4	93 \pm	92 \pm 3	98 \pm 12	97 \pm 5

Table 4 (cont.)

Binding of IFN- α 2/1 and anti-IFNAR2 mAbs to single alanine IFNAR2 mutants

mutants	% Wild IFNAR2 binding				
	IFN- α 2/1	1D3	1F3	3B7	Polyclonal
K153A	71 \pm 1	91 \pm 1	86 \pm 6	96 \pm 11	93 \pm 13
K154A	82 \pm 18	90 \pm 1	78 \pm 6	89 \pm 19	91 \pm 4
K156A	62	83 \pm 111	97 \pm 110	87 \pm 7	96 \pm 17

Experiments were carried out as described in Table 3.

Upon analysis of the hydropathy profile of the IFNAR2, 12 charged areas in clusters of 2-7 residues were selected for substitution with alanines as shown in Table 3 above. Eleven out of 12 multiple mutants expressed as immunoadhesins as detected by anti-human IgG-Fc ELISA. IFN α -2/1 exhibited no binding to IFNAR2 multiple mutants in the residues of 49-55, 68-72, 74-78, 105-109, 133-139, 153-156 while multiple alanine substitutions in the area 7-11, 29-33, 159-163, 172-173 and 187-192 exhibited no effect on the binding of IFN α -2/1. No expression was obtained from the IFNAR2 multiple mutant containing alanine substitutions in residues 145-149 of IFNAR2, indicating that residues 145-149 are important for the integrity of IFNAR2-IgG. A computer model of IFNAR2 was constructed by displaying its sequence on the backbone of tissue factor (described in Muller et al., *J. Mol. Biol.*, **256**: 144-159 (1996)). Fig. 4 shows a three dimensional rendering of this model. According to the model, residues 49-55, 68-72, 74-78, 105-109, 133-139 and 153-156, are clustered into a major area. Thus, it appears that these areas of IFNAR2 contribute to the binding of IFN- α 2/1. To confirm the integrity of these multiple substitution IFNAR2 mutants, the binding of anti-IFNAR2 mAbs to the mutants was also

determined. All the mutants expressed exhibited 35-100 % binding with at least one of these mAbs, compared to the wild type IFNAR2-IgG, indicating that these multiple mutants retain the general integrity of IFNAR2 structure. The data also demonstrated that the binding of the most potent blocking mAb 3B7 was dependent upon the residues 49-55 and 68-72 of IFNAR2, and was influenced by the residues 133-139 and 153-157. The epitopes recognized by the moderate blocking mAb 1F3 include residues 49-55, 68-72 and 74-78 in domain 1, and residues 105-109, 133-139 and 153-156 in domain 2. The critical areas recognized by mAb 1D3 are in residues 133-139 and 153-157 in domain 2 of IFNAR2 while residues 49-55 and 68-72 have some influence on mAb 1D3 binding.

To better define residues on IFNAR2 which play important roles in the binding of IFN α -2/1 and blocking mAbs, single alanine mutants were generated in the hydrophilic amino acids in residues of 49-156 (see Table 4 above). IFNAR2 single alanine mutants, D68A, E78A and W101A, exhibited no significant binding to IFN- α 2/1 while IFNAR2 single alanine mutants I104A and D105A exhibited some binding to IFN- α 2/1. Although no single residue which has crucial effects on the binding of the anti-IFNAR2 mAbs was detected, some of the multiple alanine substitution mutants shown in Table 3 exhibited a significant reduction in the binding of these mAbs. The experiments depicted in Tables 3 and 4 were done by capturing only 20-50 ng (0.2-0.5 nM) of mutant IFNAR2-IgG in goat anti-human IgG coated wells. After washing, the IFNAR2-IgG bound was detected by the addition of mAbs at a concentration of 1 μ g/ml (6 nM) or biotinylated IFN- α 2/1 at a concentration of 1 μ g/ml (50 nM). Thus, the molar ratio between

IFNAR2-IgG mutants captured and the mAb or type 1 IFN added is more than 1: 1000 fold.

Under such experimental conditions, it was possible to detect almost all IFNAR2-IgG mutant bound. Similar data were obtained from five independent experiments. In order to confirm the

findings shown in Table 3, the binding capacities of some of important IFNAR2-IgG mutants were determined using various concentration of IFN- α 2/1 and mAb 1F3 (Figs. 3A-3B). The

5 binding of IFN- α 2/1 to IFNAR2-IgG mutants D68, E78 and W101 at IFN- α 2/1 concentrations of 5-50 nM was found to be less than 10 % of binding to wild type IFNAR2-IgG while the EC 50 of

mAb 1F3 for the binding to IFNAR2-IgG mutants I104A and D105A was determined to be approximately 10 fold higher than the EC50 of mAb 1F3 for wild type IFNAR2-IgG. These

results confirmed that residues D68, E78 and W101 play a crucial role in the binding of IFN- α 2/1 and residues I104 and D105 influence the binding of IFN- α 2/1. Among all of the single mutants

tested, mutant E78A exhibited the biggest reduction (54%) in binding to mAb 1F3 as compared to wild type. The EC50 concentrations exhibited by mAb 1F3 with respect to wild type IFNAR2-

IgG, multiple 74-78 residue mutant (RSTHE(SEQ ID NO.11)/ASTAA(SEQ ID NO.12)) and

E78A mutant were determined to be 0.09 nM, 90 nM and 1.1 nM, respectively. These results

15 indicate that the residue E78 plays some role on the binding of mAb 1F3 but the cluster of R74, H77 and E78 together has a drastic effect (1000 fold reduction) on the binding of mAb 1F3 to

IFNAR2-IgG.

DISCUSSION

On the basis of structural homology, hematopoietic superfamily receptors can be divided into two classes. The first class of the hematopoietic superfamily includes receptors for human growth hormone (hGH), erythropoietin, granulocyte-macrophage colony-stimulating factors, interleukin-3, -4, -6 and -7 while the second class of the cytokine receptor family included the receptors for IFN- γ , IFN- α and IL-10. The structures of hGHR and tissue factor belong to class 1 and class 2 of cytokine receptor superfamily, respectively, and have been well characterized by mutational analysis and crystal structure analysis. The main difference in these two receptors is the angle between domains. The angle between domains in hGHR is about 85° while that of tissue factor is about 120°. The known structure of tissue factor was used as a backbone to construct a computer model of the ECD of hIFNAR2 (Figure 3). The ECD of hIFNAR2 is composed of 2 domains (approximately 100 amino acids/ domain) while the ECD of hIFNAR1 is composed of 4 domains. The multiple alanine substitutions in residues of 49-54, 68-72, 74-78, 105-109, 133-139 and 153-156 of IFNAR2 completely abolished the binding of IFN- α 2/1 as well as mAb 1F3 to IFNAR2. Upon a comparison of the computer model and the location of the residues identified in the multiple mutant analysis, it is apparent that these residues collectively form a patch. Accordingly, these residues play an important role in the binding of the ligand and mAb 1F3. Single alanine mutant analysis showed that residue D68, E78 and W101 are crucial in the binding of IFN- α 2/1 while residues I104 and D105 contribute to the binding of IFN- α 2/1. Since the residues D68, E78, W101, I104 and D105 form a small pocket in the computer model,

the data indicate that this pocket is important in binding to the ligand. Although the portion of IFNAR2 involved in the binding of mAb 1F3 exhibits a significant overlap with the portion of IFNAR2 involved in the binding of IFN- α 2/1 as determined by the multiple alanine mutation analysis, no single residue was shown to be crucial for the binding of mAb 1F3 binding whereas residues D68, E78 and W101 were shown to be crucial for the binding of IFN- α 2/1. This result indicates that the binding of blocking mAb 1F3 to the receptor is not necessarily same as the binding of ligand to the receptor.

The epitopes recognized by mAb 3B7 and mAb 1D3 were localized to domain 1 (residues 49-55 and 68-72 in particular) and domain 2 (residues 133-139 and 153-157 in particular) of IFNAR2, respectively. Although, the epitope(s) recognized by mAb 3B7 do not closely overlap with the epitope recognized by IFN- α 2/1, mAb 3B7 exhibited the most potent blocking activity. In contrast, mAb 1D3 recognized residues located in lower domain 2 of IFNAR2, which forms a part of the ligand binding areas and showed weak blocking activities in the ISGF complex formation and in anti-viral assay.

The following hybridomas have been deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD, USA (ATCC):

<u>Cell Lines</u>	<u>ATCC Accession No.</u>	<u>Deposit Date</u>
3B7		
1F3		
1D3		

These deposits were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable deposit for 30 years from the date of deposit. These cell lines will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the cell lines to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the cell lines to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC §122 and the Commissioner's rules pursuant thereto (including 37 CFR §1.14 with particular reference to 886 OG 638).

The assignee of the present application has agreed that if the deposited cell lines should be lost or destroyed when cultivated under suitable conditions, they will be promptly replaced on notification with a specimen of the same cell line. Availability of the deposited cell lines is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.